

# CURRICULUM VITAE

## Fabio Happacher

Laboratori Nazionali di Frascati dell'INFN  
fabio.happacher@lnf.infn.it

### Personal Information

### Education

- Università dell'Aquila (L'Aquila, Italy) - Department of Physics, Doctorate in Physics, February 16, 2001. The argument of the thesis is: "Study of leptonic decays of hadrons with heavy flavor quarks produced in  $p\bar{p}$  interactions at  $\sqrt{s} = 1.8$  TeV".
- Università di Roma "Tor Vergata" (Rome, Italy), *Laurea* in Physics, Nuclear and Subnuclear Physics, December 19, 1996. The argument of the thesis is: "Study of  $W$  boson production in association with jets containing heavy flavor quarks and measurement of  $t\bar{t}$  pairs production cross section".
- Liceo Scientifico Statale di Grottaferrata "Bruno Touschek".  
*Maturità scientifica*, July 1989.

### Professional Experience and Charges

- Group Leader of the LNF-INFN group working on the Mu2E experiment at Fermilab
- Achieved the National Scientific Qualification (ASN) of the Education Ministry for Associate Professor, Jan 2014
- Elected Member of the Users Executive Committee at Fermilab (UEC) -Sep 2014 to date. The purpose of the Committee is to provide a forum for discussion of scientific and administrative matters relevant to the organization and functions of the Laboratory. One of the most important activity is the annual visit to Washington DC congressional offices and speak with members of the United States Senate and House of Representatives seeking support for HEP.
- Staff Research Scientist, Laboratori Nazionali di Frascati of INFN (Italy), October 2004 - to date.

→ Mu2e collaboration (Fermilab) - Currently, L3 manager for the mechanics design and integration of the calorimeter. I coordinate the Mechanics and Electronics engineering team that is designing all the components of the calorimeter giving them all the physics inputs to achieve the best design performances. As of spring 2016, in view of the CD-3 approval review, most of the calorimeter design is final and we are building real scale prototypes of all the critical parts to optimize their design or spot critical issues.

Characterization, simulations, design and realizations and tests of small size prototypes for the crystals calorimeter readout by new generation solid state photosensors for the Mu2E experiment. Responsible for the design and assembly of the prototypes; three matrices made of different crystal types and equipped with photosensors, FEE electronics and LED/Laser calibration systems have been extensively tested in electron and photons Test Beam campaigns at the Frascati LNF Beam Test Facility and at Mainzer Microtron (Mainz) respectively.

→ KLOE-2 collaboration (LNF) - Realization and characterization of a new forward calorimeter prototype surrounding the inner quadrupoles, QCALT, devoted to the identification of photons from rare  $K_L$  decays.

→ KLOE-2 collaboration (LNF) - Responsible for the design, prototyping and realization of a small angle crystal calorimeter to improve the acceptance for tagging low energy photons coming from rare Eta and Kaon event decays ( $K_s \rightarrow \gamma\gamma, \eta \rightarrow \pi_0\gamma\gamma$ ).

→ CDF II Experiment - Study of the sub-sample of dimuon events that appears to be responsible for the observed discrepancies of the  $b$  quark pairs cross section measurements,  $\sigma_{b\bar{b}}$ , with respect to the prediction;

→ CDF II Experiment -  $b$ -quark pairs production cross section measurement,  $\sigma_{b\bar{b}}$ , looking at events with muon pairs with large impact parameter originating from semileptonic  $b$ -quark decays;

→ CDF II Experiment - Study of  $B$  into  $J/\Psi$  decays, inclusive  $b$ -quark production cross section measurement via  $B^+ \rightarrow J/\Psi K^+$  channel;

→ CDF II Experiment - Responsible for the energy scale calibration and maintenance of the CDF central hadron calorimeters;

→ CDF II internal referee for  $Z'$  and high mass dielectron searches.

→ KLONE collaboration (LNF) - Participation to the R&D project for the measurement of the neutron detection efficiency of a KLOE electromagnetic calorimeter prototype (Lead-scintillating fibers). Three test-beams are carried on at the Svedberg Laboratory neutron beam facility at Uppsala varying neutron beam energy and intensity.

- Post doctoral Researcher (*Assegno di Ricerca*), Laboratori Nazionali di Frascati of INFN (Italy), February 2002 to September 2004. The research activity is carried on at the Collider Detector at Fermilab experiment (CDF II).

- Development of solutions to compensate for the reduced charge integration time of the CDF central calorimeter photomultipliers with 400 ns bunch spacing of the upgraded Tevatron;

- Development of a real-time object oriented software package for CDF to analyze laser data;

- Responsible for the installation, debug and performance monitoring of the new discriminators of the electromagnetic and hadronic calorimeters and for the calibration of the timing system;

- Physics Research Doctorate, L'Aquila University (Italy), November 1997 - March 2001. The research activity is carried on at the Collider Detector at Fermilab experiment (CDF).

- Study of the heavy flavor content of events containing a low- $p_T$  lepton plus jets; CDF Run I data show an excess of events with kinematic properties that are consistent with the production of a light scalar quark;

- Search for narrow bound resonances in the dimuon sample with mass between the  $J/\Psi$  and the  $\Upsilon$ ;

- Participation to the Stanford Linear Accelerator Laboratory Summer School: Secrets of B Mesons, August 2002.

- Fermilab - Guest Scientist at CDF.  
February to September 1997 - April to December 2001.

- Tor Vergata University (Rome) and Laboratori Nazionali di Frascati of INFN, experimental *Laurea* thesis carried on at the Collider Detector at Fermilab experiment (CDF), June 1995 - December 1996.

- Revision of the  $b$ -quark tagging efficiency measurement leading to the new and final Run I top pairs production cross section in the  $W \rightarrow e(\mu), \nu_e(\nu_\mu) + \text{jets}$  (lepton plus jets) channel. Development of the energy corrections for  $b$  flavoured jets to improve the top events acceptance and mass resolution.

- Pre graduate Fellow at Fermi National Accelerator Laboratory (Fermilab), July 1995 to October 1995. He works in the CDF experiment and gains the first particle physics experience during the Thesis activity.

## Outreach

- Responsible for educational experiments “Revealing particles with calorimeters” at the LNF “Incontri di Fisica: refresher/educational course for high school teachers” since year 2001.
- Responsible for tutoring undergraduate students at the LNF Summer Stages.
- Responsible for tutoring undergraduate students at the Fermilab Summer Student Program.

## Participation at Conferences

- The Mu2e Experiment at Fermilab LASNPA -WONP-NURT 2017 (Latin-American Symposium on Nuclear Physics and Applications), La Habana (Cuba) 2017, October 23-27
- The Mu2E experimet at Fermilab  
New trends in High Energy Physics, Budva, Becici, (Montenegro), 2016, October 2-8
- The Mu2E crystal calorimeter  
INSTR17: Instrumentation for Colliding Beam Physics, Novosibirsk (Russia) 2016, February 27- March 3
- L'Esperimento Mu2e al Fermilab; 101° Congresso della Società Italiana di Fisica, Rome, 2015, September 21-25.
- A Crystal Calorimeter for the Mu2e Experiment; CHEF 2013 - Calorimetry for High Energy Frontier, Paris (France) - 2013, April 22 - 25.
- CCALT: crystal calorimeter at KLOE-2; 14<sup>th</sup> International Conference on Calorimetry in High Energy Physics, Beijing (China) - 2010, May 10 - 14.
- Study of Multimuons Events at CDF; Les Rencontres de Physique de la Valle d'Aoste 2009, La Thuile, Aosta Valley (Italy) - 2009, March 1-7.
- Study of Multimuons Events at CDF; XVII International Workshop on Deep-Inelastic Scattering and Related Subjects DIS 2009, Madrid (Spain) - 2009, April 26-30.
- Study of Multimuons Events at CDF; The 2009 Europhysics Conference on High Energy Physics, Krakow (Poland) - 2009, July 16-22.
- Measurement and simulation of the neutron response and detection efficiency of a Pb-scintillating fiber calorimeter; SCINT 2007, IEEE-9th International conference on Inorganig Scintillators and their Applications, Winston-Salem, NC USA - 2007, June 4-8.
- Status of the observed and predicted  $b\bar{b}$  production at the Tevatron; XIV International Workshop on Deep Inelastic Scattering, Tsukuba (Japan) - 2006, April 20-24.
- High Energy Physics and its Experiments; WYP-2005, World Year of Physics dissemination program, Seminar at Liceo Serpieri - Rimini (Italy) 2005, April 15.
- Ecfa Study, First workshop on Physics and Detectors for a Linear Collider, Montpellier (France) - 2003, November 13-16.
- Properties of heavy flavour jets produced at Tevatron  $\sqrt{s} = 1.8$  Tev, Incontri di

Fisica delle Alte Energie, XV (IFAE), Lecce (Italy) - 2003, April 23-26.

- Higgs searches at Tevatron, Tor Vergata University, Rome (Italy) - November 2000.
- The latest Run I result of the top cross section measurement, New Perspectives '99, Fermilab - June 1999.

## Research Activity details

- Current position

Mu2e The Mu2e experiment looks for the CLFV decay  $\mu \rightarrow e$ . It requires a high energy and time resolution calorimeter to complement the tracking detector in the measurement of track energy to separate conversion electrons from  $\mu \rightarrow e$ , with 105 MeV/c momentum, from the electrons coming from the decay in orbit of orbiting muons in the Al target,  $\mu \rightarrow e \nu$ , that have a momentum spectrum with 105 MeV/c endpoint. It also needs to reject background due to reconstruction errors or cosmic ray muons. The requirements for the calorimeter are 5% energy resolution,  $<1$  ns time resolution and  $\sim$  cm spatial resolution. In the 100 MeV energy region a total absorption calorimeter employing a continuous and homogenous medium is needed. The Mu2e collaboration has chosen to consider a crystal calorimeter made of two possible type of scintillating crystals, LYSO or BAF<sub>2</sub>. Given the experience acquired in crystal calorimetry for the KLOE-2 collaboration the Frascati group has been asked to take the responsibility of designing and realizing the calorimeter detector.

I personally followed all the stages of the Mu2e approval by the funding agencies in USA and in Italy. We showed all the phases of the R&D we have been carrying on to study the performances and to tune them with the requirements highlighted by simulation studies.

We built different crystals array prototypes ameliorating everytime the FEE electronics and the photosensors control. All these prototypes have been tested with electrons or photons beams at BTF (Frascati) and Mami (Mainz).

The calorimeter design is currently described in all its details in the Mu2e Technical Design Report [arXiv:1501.05241](https://arxiv.org/abs/1501.05241) prepared for the CD-3 approval expected by spring 2016. **Nucl. Instrum. Meth. A 718, 56 (2013)** describes the Mu2e calorimeter project together with the energy resolution results of the first test beam at Mami.

As L3 manager I'm currently responsible for the mechanics design an integration of the calorimeter. The electron energy and timing resolutions optimization drive the technical choices of all the components of the calorimeter: type of crystal, photosensor, FEE electronics, digitizations; all these elements need to be engineered and be housed in the calorimeter. I have the responsibility of transferring all the physics inputs to the engineer team that is designing the mechanics of the calorimeter and to play an interface role with the Muon Beamline group that is integrating all the detectors in the Detector Solenoid.

The calorimeter consists of two identical annuli 1500 mm outer diameter, 700 mm inner diameter, composed each of an inner cylinder made of light composite material, an outer stainless steel structural ring, one light frontal plate, a stack of 800 self-standing crystals equipped with two SiPM photosensors and FEE

electronics. It is also equipped with a sophisticated cooling system to extract the electronics dissipated power and to cool down the photosensors in order to recover from deterioration of response due to an increased leakage currents as effect of radiation damages.

CDF The measurement of  $\sigma_{b\bar{b}}$  described in **Physical Review D 77 (2008) 072004** agrees with NLO predictions. A detailed investigation of the dimuon data sample indicates that the discrepancies with respect to previous measurement are related to the requirement that the two muons tracks are reconstructed with the silicon detector with hits in the two inner layer and at least two in the four outer layers; in other words, that at least one of the two muons has to decay inside the 1.5 cm radius beam pipe. When this silicon requirement is released the measurement agrees with previous measurements and does not match the theory.

This is due to a set of events with interesting properties that were unaccounted for when using looser silicon cuts. A large fraction of these events is attributed to muons arising from in flight decays and punch-throughs of hadronic prongs of  $K_s^0$ . The remaining fraction consists of events where the muons show a very long impact parameter tail and have a very high muon multiplicity in a narrow cone around the primary muons. These events cannot be explained in terms of SM events or detector failures. The analysis is published in **Eur. Phys. J. C 68, 109**.

CDF At the Tevatron the measurement of the bottom quark cross section,  $\sigma_b$ , and the measurement of the cross section for producing both  $b$  and  $\bar{b}$  quarks centrally and above a given momentum thresholds,  $\sigma_{b\bar{b}}$ , is an important test of the power of QCD to predict absolute rates in hadronic collisions. In QCD calculations, the long- and short- distance dynamics of the hadronic hard-scattering cross section are factorized into nonperturbative parton distribution functions (PDF) and fragmentation functions, and perturbatively calculable hard-scattering functions.

In **Physical Review D 73 (2006) 014026** all the Tevatron measurements have been compared to the same QCD prediction and they appear to be inconsistent among themselves and higher than the expectation.

Measurements of inclusive  $b$  cross sections,  $\sigma_b$ , and  $b$  pairs cross sections,  $\sigma_{b\bar{b}}$ , are sensitive to different contributions of the perturbative level of the QCD hard scattering calculations; inclusive  $b$  productions is contributed by Next to Leading Order terms (NLO)  $b\bar{b}$  pairs production is dominated by leading order terms (LO). Precise measurements can contribute to assess the role of different terms in the perturbative prediction and improve the modeling of fragmentation functions.

The analysis described in **Physical Review D 75 (2007) 012010** measures the  $\sigma_b$  cross section using the decay  $B^\pm \rightarrow J/\Psi K^\pm$  with  $J/\Psi \rightarrow \mu^+ \mu^-$  reconstructing the invariant mass peak of the decay particles in different  $p_T$  bins.

Acceptancies are measured from Monte Carlo using a NLO calculation in conjunction with EVTGEN for  $B$  hadrons decays and the GEANT based simulation of the CDF detector. To rely on Monte Carlo the the most relevant efficiencies and acceptancies are cross-checked using data control sample.

A second analysis determines the production rate of muons originating from  $b$ - and  $\bar{b}$ -quarks semileptonic decays. A fit to the impact parameter distributions of muons allows to distinguish from muons coming from  $b$ -  $c$ -quark decays or prompt decays of quarkonia and Drell-Yan production. The fit uses heavy flavor impact parameter templates determined using the HERWIG Monte Carlo generator. Since the analysis is based on lifetime distributions, HERWIG has to reproduce the measured abundances of heavy flavor hadrons.

To have the right  $b$ - and  $c$ - hadrons composition, produced by the heavy quarks fragmentation, HERWIG has been tuned to reproduce the PDG fragmentation fractions for  $b$ -quark and the experimental results from BELLE, CLEO, HERA, LEP for  $c$ -quark. This work is documented in **Physical Review D 77 (2008) 072004**.

Both results agree with the Standard Model prediction.

CDF Responsible for the absolute energy scale calibration of the Central (CHA) and EndWall (WHA) hadron calorimeters and for the maintainance of the calorimeters hardware during the data-taking operations. The energy scale is set using a mixture of techniques: the photomultipliers gain stability is monitored by means of frequent nitrogen Laser runs. The laser light shines the light guides facing the photomultiplier photocatodes thus originating a reference pulse height monitored weekly. The scintillator output degradation is checked looking at real energy deposition in the calorimeter towers using minimum ionizing particles (muons from  $J/\Psi$  decays reconstructed using tracking information). For this purpose a C++ based package is developed to run automatically every  $50 \text{ pb}^{-1}$  of integrated luminosity on the dimuon trigger data set. The hadronic energy response to muon of each calorimetric tower is compared to the RUN I response (when the calorimeter was calibrated using  $^{137}\text{Cs}$  source system). The absolute energy scale is determined at the level of 1-2% precision.

A  $^{137}\text{Cs}$  sources system determines the absolute energy scale for the WHA calorimeter; this procedure relies on the test beam of '83 - '85 that used  $\pi$ 's with energy greater than 10 GeV.

The absolute energy scale calibration and its stability contributes to the improvement of the jet energy scale described in **Nucl. Instrum. Meth. A 566 (2006) 375**. This is very important in reducing the sistematic uncertainty in precise elettroweak measurements like the top quark mass.

KLONE The detection of neutrons is usually performed with organic scintillators where the neutrons undergo elastic scattering with proton of hydrogen atoms. Nevertheless rem counters also take advantage of medium-high  $Z$  materials to increase

the detection response through inelastic processes that produce secondary neutrons. The KLONE collaboration performed three test beams, at the neutron facility of Uppsala Svedberg Laboratory, using the KLOE electromagnetic prototype. This calorimeter is composed of 200 layers of 1 mm blue scintillating fibers glued in grooved lead layers. The trigger/DAQ system is performed with NIM/VME technology. The test beam measurements show indeed an enhancement of the neutron efficiency and are in fair agreement with the FLUKA simulation.

The detector is calibrated using cosmic ray runs. The contribution of a beam halo to the detection efficiency is determined studying the time of flight distributions. A NE-110 scintillator detector is built for reference measurements, read-out with standard photomultipliers. Design and construction of segmented WLS fibers - scintillator strips detector for determination of the geometrical beam profile, read-out with multi-anode photomultipliers. This measurement is presented at the **IX IEEE Scint07 Conference**. The measured efficiency ranges between 30 to 50%, depending on energy, at the lowest trigger threshold; four times larger than expected for an equivalent scintillator thickness.

KLOE-2 The KLOE-2 proposal includes the addition of a new forward calorimeter, QCALT, to increase the photons detection acceptance in rare events decays ( $K_L \rightarrow 2\pi^0$ ). QCALT is a high granularity calorimeter based on tungsten scintillator tiles + WLS fibers read out with silicon photodevices (SIPM). The calorimeter consists of around 2000  $5 \times 5 \times 5$  cm<sup>3</sup> tiles, assembled in two barrel calorimeters 12 wedges each, for 1 m length and 5 cm thickness.

The R&D phase consisted on

- characterization of the SIPM response with a pulsed LED;
- study of the optical connection of the SIPM to scintillating fibers and of the response to a  $\beta$  source;
- construction of the first prototype cell, instrumented with Hamamatsu MPPC of 400 pixels and testing with cosmic rays and the electron beam at the Beam Test Facility of LNF.

Development of a DAQ/trigger system to read out the SIPM signals. The detector is fully operational in the KLOE2 apparatus.

KLOE-2 In the new design of the Dafne interaction region the position of the quadrupoles increases the acceptance of the central calorimeter from 21 to 18 degrees. In this volume is possible to insert few crystals to improve the acceptance for tagging low energy photons coming from rare Eta and Kaon event decays ( $K_s \rightarrow \gamma\gamma, \eta \rightarrow \pi_0\gamma\gamma$ ).

The proposed solution is to insert an homogeneous calorimeter immersed in a 0.52 kgauss axial magnetic field, CCALT, based on the new generation of crystals, LYSO, characterized by a very high light yield (60% of NaI), 40 ns emission

time, high density and radiation length. These crystals match the request for a high efficiency for low energy photons and an excellent time resolution needed by the calorimeter to sustain the high level of machine background events.

CCALT needs to maximize the light collection to improve its timing performance so that we selected as photosensor the fastest APD on the market from Hamamatsu.

We built a prototype matrix of  $2 \times 2 \times 15 \text{ cm}^3$  LYSO crystals read-out by  $0.5 \times 0.5$  Hamamatsu avalanche photodiodes (APD) and we tested it using cosmic rays at the electrons beam test facility of Frascati.

The measurements show a very good time resolution of around  $200 \text{ ps}$  and a light yield of 2000 photoelectrons per MeV.

We studied the energy resolution of this kind of calorimeters, that is limited by the containment of the shower and by the electronic noise. For this purpose we built a set of much larger prototypes, the first consisting of 21 crystals, 9 LYSO for an inner matrix and 12 PBWO for outer leakage recovery (3 Moliere radius in total), we measured 5% energy resolution @100 MeV.

The calorimeter is composed of four aluminum wedges per side; each wedge has three sectors with a granularity of four projective crystal. Every crystal is readout with large area Sipm.

I followed the design and the realization of the mechanical shells with a characteristic shape of truncated pyramid with a dodecagonal base manufactured using Electro Discharge Machining; the design and realization of the Sipm holder using a 3D PVC printer, the international tender for the purchase of the crystals and the photosensors. After a phase of quality assurance of all the components, all the parts were assembled and tested as a whole. The calorimeter is installed over the inner quadrupoles of Kloe and is fully operational as described in **Acta Phys. Polon. B 46, no. 1, 87 (2015)**.

- Post Doctoral Research

The LNF group is historically responsible for the building, upgrading, maintenance and calibration of the central hadronic calorimeter of the CDF experiment leading the commissioning of the central hadron calorimeters for Run II.

The hadron calorimeters underwent minor upgrades going from Run I to Run II; the front end electronics has been replaced to sustain the higher luminosity acquisition rate; all the photomultipliers functionality has been restored. The shorter acquisition gate causes a loss of the charge integration of photomultipliers pulses and various techniques are developed to set back and monitor the absolute energy scale of the calorimeters using also the first collision data from the Tevatron.

An object oriented software package is developed to analyze (online) laser runs for the calibration of the electronic chain from the photomultipliers to the front-end electronics.

Involved in the installation, calibration, performance monitoring and commissioning of the new timing system for the hadronic and electromagnetic calorimeters (**Nucl. Instrum. Meth. A 565 (2006) 538**).

The timing calibrations of the detectors, the debugging of the new discriminators installation and their thresholds settings have been checked through many “engineering” runs of  $p\bar{p}$  collisions that have been analyzed using software packages developed in C++ language within the AC++ framework of the CDF offline software.

- Physics Research Doctorate

Detailed study of the data set containing a low- $p_T$  lepton plus jets with heavy flavor searching for a light scalar quark (mass in the range of  $\sim 3.5$  GeV/ $c^2$ ) with a long lifetime and an anomalous semileptonic branching ratio. At the Tevatron energies, heavy flavor jets coming from  $b$  and  $c$  quarks hadronization show a displaced vertex with respect to the primary vertex of the event due to the effect of the long lifetime of heavy flavor hadrons ( $\tau \simeq 1$  ps) together with the Lorentz boost acquired by jets. These heavy flavor jets are identified using tagging algorithms (SECVTX, Jetprobability) based on the location of the secondary vertex. The semileptonic decay rate of heavy flavor quarks is instead measured looking for soft (low momentum) leptons not isolated in the calorimeter, contained in jets with secondary vertices. These leptons are found by the Soft Lepton Tagging algorithm (SLT).

The basic results of this analysis is the observation of an excess of events containing an additional soft-lepton inside one of the jets recoiling against the jet containing the trigger lepton significantly larger than what is expected for the production and semileptonic decay of  $b$  and  $c$  hadrons. This part of the analysis has been published in **Physical Review D. D69:072004,2004**.

Also an excess of events with an additional lepton inside the jet containing the trigger lepton was observed. These events and their kinematical properties are described in the paper published in **Physical Review D72:072002,2005**.

As a working hypothesis, the excess can be interpreted (in terms of supersymmetric signal) as the pair production and decay of a spin 0 scalar quark. The best candidate is the supersymmetric partner of the  $b$  quark:  $\tilde{b} \rightarrow c\tilde{\chi}^\pm$  and  $\tilde{\chi}^\pm \rightarrow \tilde{l}l$ , with  $m_{\tilde{b}} = 3.65$  GeV/ $c^2$ ,  $m_{\tilde{\nu}} = 0$  and  $\tau_{\tilde{b}} \simeq 1$  ps. This process has been implemented in the Herwig Monte Carlo generator using a matrix element calculation by K. Hikasa and M. Kobayashi for SUSY models and reproduces most of the features of the observed anomalies. These results are explained in terms of SUSY signal in **Physical Review D73:014025,2006**.

Based on the interpretation of a scalar quark, this analysis was extended to a search of narrow bound states of two scalar quarks by studying the dimuon invariant mass spectrum between 6 and 9 GeV/ $c^2$ . This latter study is published in **Physical Review D72:092003,2005**.

- Laurea Thesis

The argument of the thesis is the study of the  $W + \text{jet}$  sample where  $t\bar{t}$  production

events can be isolated and top production cross section measured. Each top quark in the event decays to a  $W$  boson and a  $b$  quark with a Branching Ratio of  $\sim 100\%$  and there are different research channels according to the decay of the two  $W$  bosons present in the event (to hadrons or to leptons). It is used the “lepton plus jets” channel, where one of the two  $W$  decays to leptons,  $W \rightarrow l\nu_l$  ( $l = e, \mu$ ) and the other to two adronic jets. The event topology consists of a lepton with high momentum isolated in the calorimeter, an unbalance of transverse energy in the calorimeters ( $\cancel{E}_T$ ) indicating the presence of an undetected neutrino and at least four jets; furthermore it is required that at least one of the jets contains a secondary vertex, reconstructed by the SECVTX algorithm, in order to enhance the presence of  $b$  quarks.

The measurement of the top production cross section at Tevatron provides the opportunity to test the Standard Model predictions comparing the data with the QCD +  $t\bar{t}$  prediction of  $W$ + jets events as a function of the multiplicity and transverse momentum of the jets in the event.

The highest contribution to background events for top comes from the direct QCD production of  $W$  boson associated to jets from gluons radiation that split to heavy flavor pairs. The amount of these events is estimated from Monte Carlo.

The excess over the background of events with at least three jets is attributed to top; geometrical acceptancies of the analysis cuts and tagging efficiencies are measured in a top Monte Carlo sample. With an integrated luminosity of  $\int \mathcal{L} dt = 105 \pm 4 \text{ pb}^{-1}$  we count 29  $W + \geq 3$  jets events with  $7.99 \pm 0.98$  background events, corresponding to a  $\sigma_{t\bar{t}} = 5.1 \pm 1.5 \text{ pb}$ . This value is the official result of the CDF Run I and is published on **Physical Review D64:032002,2001** .

## List of Papers

### Primary Publications

- "A LYSO crystal calorimeter for the Mu2e experiment".  
Proceedings of the CHEF2013 Conference - Eds. J.-C. Brient, R. Salerno, and Y. Sirois  
ISBN number 978-2-7302-1624-1
- "Measurement of correlated b-bbar production in p-pbar collisions at  $s^{**}(1/2) = 1960$  GeV," Phys. Rev. D **77** (2008) 072004.
- T. Aaltonen *et al.* [CDF Collaboration], "Study of multi-muon events produced in  $p\bar{p}$  interactions at  $\sqrt{s} = 1.96$  TeV," Eur. Phys. J. C **68**, 109 (2010).
- "Measurement of the  $B^+$  production cross section in  $p\bar{p}$  collisions at  $\sqrt{s} = 1960$  GeV", Phys. Rev. D **75** 2007012010.
- F. Happacher, "Measurement and simulation of the neutron response and detection efficiency of a Pb-scintillating fiber calorimeter", in proceedings of IX IEEE Scint07 Conference - Winston Salem, North Carolina USA, june 4-8, 2007.
- F. Happacher, "Status of the observed and predicted  $b\bar{b}$  production at the Tevatron", in proceedings of XIV International Workshop on Deep Inelastic Scattering - Tsukuba (Japan), april 20-24, 2006.
- F. Happacher, "Properties of jets with heavy flavor produced in in p anti-p interactions at  $\sqrt{s} = 1.8$ -TeV", in proceedings of IFAE 2003 - Lecce, april 23-26, 2003.
- F. Happacher, P. Giromini and F. Ptohos, "Status of the observed and predicted b barb production at the Tevatron," Phys. Rev. D **73** (2006) 014026.
- G. Apollinari *et al.*, "Search for narrow resonances below the Upsilon mesons," Phys. Rev. D **72** (2005) 092003.
- G. Apollinari *et al.*, "Phenomenological study of the atypical heavy flavor production observed at the Fermilab Tevatron," Phys. Rev. D **73** (2006) 014025
- G. Apollinari, I. Fiori, P. Giromini, F. Happacher, S. Miscetti, A. Parri and F. Ptohos, "Study of sequential semileptonic decays of b hadrons produced at the Tevatron," Phys. Rev. D **72** (2005) 072002.
- Heavy flavor properties of jets produced in p anti-p interactions at  $\sqrt{s} = 1.8$  TeV.  
Nov 2003, D. Acosta *et al.*, Published in Phys. Rev. D **69** (2004) 072004
- Study of the heavy flavor content of jets produced in association with  $W$  boson in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.8$  TeV.  
Sep 2001, D. Acosta *et al.*, Published in Phys. Rev. D **65** (2002) 052007

- Measurement of the  $t$  anti- $t$  production cross-section in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.8$  TeV.  
Jan 2001, T. Affolder *et al.*, Published in Phys. Rev. D **64** (2001) 032002.
- Cordelli:2013mka M. Cordelli, E. Dan, S. Giovannella, M. Gatta, F. Happacher, A. Luc, S. Miscetti and A. Saputi *et al.*, “CCALT: A Crystal CALorimeter with Timing for the KLOE-2 upgrade,” Nucl. Instrum. Meth. A **718**, 81 (2013).

### Internal CDF Publications

1. CDF9230, PRD DRAFT: Study of multi-muon events produced in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV - 20/10/08;
2. CDF8733: Measurement of the correlated  $b\bar{b}$  production - 11/6/07;
3. CDF8394: Measurement of the  $B^+$  production cross section at  $\sqrt{s}=1.96$  TeV - 21/7/06;
4. CDF8507: Tuning of herwig version 6.5 - 25/9/06;
5. CDF7543, NIM Draft: Determination of the jet Energy scale at the Collider Detector at Fermilab - 11/7/05;
6. CDF7599: Cross-check of the CHA and WHA calibrations - 26/4/05;
7. CDF6891: Energy scale calibration of CDF Hadron Calorimeters - 23/2/04;
8. PRD draft: Study of the semileptonic decays of  $b$  hadrons produced at Tevatron - 2003;
9. PRD draft: Search for narrow resonances below the  $\Upsilon$  mesons - 2003;
10. PRD draft: A possible explanation for the anomalous heavy flavor production observed at Tevatron - 2003;
11. CDF5856: Timing calibration of CDF hadron calorimeters - 14/2/02;
12. CDF4795: Observation of semileptonic decays of heavy quarks produced in  $p\bar{p}$  interactions and the existence of one scalar quark - 29/3/2000;
13. CDF4257: Measurement of the data to simulation scale factor for the tagging efficiency of SECVTX and JPB - 28/3/99;
14. CDF4303: Method II measurement of the top cross section with W+3,4 jet events with SECVTX, JPB and SLT tags - 26/3/99;
15. CDF4860: Measurement of the dimuon invariant mass between the Psi and the Upsilon - 25/1/1999;
16. CDF4115: Heavy flavor content of jets in the data and in Herwig simulations - 19/3/97;

17. CDF3855: Measurement of the  $t\bar{t}$  production cross section using SECVTX tags - 11/2/97;
18. CDF3861: Measurement of the  $t\bar{t}$  production cross section using SLT tags - 4/9/96;
19. CDF3876: Measurement of the  $t\bar{t}$  production cross section using Jetprobability tags - 11/9/96;
20. CDF3485: Evidence for a b-bbar state of mass 109 GeV - 17/1/96;
21. CDF3457: ZZJSCO: jet corrections for the study of ttbar, VV and VH production - 19/12/96;
22. CDF3602: Study of the top invariant mass distribution using  $W + \geq 4$  jet events - 11/3/96;
23. CDF3610: Study of the invariant mass distribution of the two highest  $E_t$  jets in  $V + \geq 2$  jet events with a SECVTX tag - 15/3/96.

# Publications of Fabio Happacher

1. A. Anastasi *et al.* [KLOE-2 Collaboration], “Limit on the production of a new vector boson in  $e^+e^- \rightarrow U\gamma$ ,  $U \rightarrow \pi^+\pi^-$  with the KLOE experiment,” *Phys. Lett. B* **757**, 356 (2016) doi:10.1016/j.physletb.2016.04.019 [arXiv:1603.06086 [hep-ex]].
2. T. A. Aaltonen *et al.* [CDF Collaboration], “Measurement of the forward-backward asymmetry of top-quark and antiquark pairs using the full CDF Run II data set,” Submitted to: *Phys.Rev.D* [arXiv:1602.09015 [hep-ex]].
3. A. Anastasi *et al.* [KLOE-2 Collaboration], “Precision measurement of the  $\eta \rightarrow \pi^+\pi^-\pi^0$  Dalitz plot distribution with the KLOE detector,” arXiv:1601.06985 [hep-ex].
4. A. Anastasi *et al.* [KLOE-2 Collaboration], “Measurement of the  $\phi \rightarrow \pi^0 e^+ e^-$  transition form factor with the KLOE detector,” *Phys. Lett. B* **757**, 362 (2016) doi:10.1016/j.physletb.2016.04.015 [arXiv:1601.06565 [hep-ex]].
5. T. A. Aaltonen *et al.* [CDF Collaboration], “Measurement of the Forward-Backward Asymmetry in Low-Mass Bottom-Quark Pairs Produced in Proton-Antiproton Collisions,” Submitted to: *Phys.Rev.D* [arXiv:1601.06526 [hep-ex]].
6. T. A. Aaltonen *et al.* [CDF Collaboration], “Measurement of the  $B_c^\pm$  Production Cross Section in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96$  TeV,” *Phys. Rev. D* **93**, no. 5, 052001 (2016) doi:10.1103/PhysRevD.93.052001 [arXiv:1601.03819 [hep-ex]].
7. T. A. Aaltonen *et al.* [CDF Collaboration], “Search for a Low-Mass Neutral Higgs Boson with Suppressed Couplings to Fermions Using Events with Multiphoton Final States,” Submitted to: *Phys.Rev.Lett.* [arXiv:1601.00401 [hep-ex]].
8. N. Atanov *et al.* [Mu2e Collaboration], “Measurement of time resolution of the Mu2e LYSO calorimeter prototype,” *Nucl. Instrum. Meth. A* **812**, 104 (2016) doi:10.1016/j.nima.2015.12.055 [arXiv:1509.04468 [physics.ins-det]].
9. A. Anastasi *et al.*, “Limit on the production of a low-mass vector boson in  $e^+e^- \rightarrow U\gamma$ ,  $U \rightarrow e^+e^-$  with the KLOE experiment,” *Phys. Lett. B* **750**, 633 (2015) doi:10.1016/j.physletb.2015.10.003 [arXiv:1509.00740 [hep-ex]].
10. T. A. Aaltonen *et al.* [CDF Collaboration], “Measurement of vector boson plus  $D^*(2010)^+$  meson production in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV,” *Phys. Rev. D* **93**, no. 5, 052012 (2016) doi:10.1103/PhysRevD.93.052012 [arXiv:1508.06980 [hep-ex]].
11. T. A. Aaltonen *et al.* [CDF Collaboration], “Study of the energy dependence of the underlying event in proton-antiproton collisions,” *Phys. Rev. D* **92**, no. 9, 092009 (2015) doi:10.1103/PhysRevD.92.092009 [arXiv:1508.05340 [hep-ex]].

12. T. A. Aaltonen *et al.* [CDF Collaboration], “Measurement of the Production and Differential Cross Sections of  $W^+W^-$  Bosons in Association with Jets in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96$  TeV,” *Phys. Rev. D* **91**, no. 11, 111101 (2015) Addendum: [*Phys. Rev. D* **92**, no. 3, 039901 (2015)] doi:10.1103/PhysRevD.91.111101, 10.1103/PhysRevD.92.039901 [arXiv:1505.00801 [hep-ex]].
13. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the top-quark mass in the  $t\bar{t}$  dilepton channel using the full CDF Run II data set,” *Phys. Rev. D* **92**, no. 3, 032003 (2015) doi:10.1103/PhysRevD.92.032003 [arXiv:1505.00500 [hep-ex]].
14. T. A. Aaltonen *et al.* [CDF Collaboration], “First measurement of the forward-backward asymmetry in bottom-quark pair production at high mass,” *Phys. Rev. D* **92**, no. 3, 032006 (2015) doi:10.1103/PhysRevD.92.032006 [arXiv:1504.06888 [hep-ex]].
15. T. A. Aaltonen *et al.* [CDF Collaboration], “Search for Resonances Decaying to Top and Bottom Quarks with the CDF Experiment,” *Phys. Rev. Lett.* **115**, no. 6, 061801 (2015) doi:10.1103/PhysRevLett.115.061801 [arXiv:1504.01536 [hep-ex]].
16. T. A. Aaltonen *et al.* [CDF and D0 Collaborations], “Tevatron Combination of Single-Top-Quark Cross Sections and Determination of the Magnitude of the Cabibbo-Kobayashi-Maskawa Matrix Element  $V_{tb}$ ,” *Phys. Rev. Lett.* **115**, no. 15, 152003 (2015) doi:10.1103/PhysRevLett.115.152003 [arXiv:1503.05027 [hep-ex]].
17. G. Pezzullo *et al.*, “Progress Status for the Mu2e Calorimeter System,” *J. Phys. Conf. Ser.* **587**, no. 1, 012047 (2015). doi:10.1088/1742-6596/587/1/012047
18. T. A. Aaltonen *et al.* [CDF Collaboration], “Measurement of central exclusive  $\pi^+\pi^-$  production in  $p\bar{p}$  collisions at  $\sqrt{s} = 0.9$  and 1.96 TeV at CDF,” *Phys. Rev. D* **91**, no. 9, 091101 (2015) doi:10.1103/PhysRevD.91.091101 [arXiv:1502.01391 [hep-ex]].
19. T. Aaltonen *et al.* [CDF and D0 Collaborations], “Tevatron Constraints on Models of the Higgs Boson with Exotic Spin and Parity Using Decays to Bottom-Antibottom Quark Pairs,” *Phys. Rev. Lett.* **114**, no. 15, 151802 (2015) doi:10.1103/PhysRevLett.114.151802 [arXiv:1502.00967 [hep-ex]].
20. A. Anastasi *et al.* [KLOE-2 Collaboration], “Search for dark Higgsstrahlung in  $e^+e^- \rightarrow \mu^+\mu^-$  and missing energy events with the KLOE experiment,” *Phys. Lett. B* **747**, 365 (2015) doi:10.1016/j.physletb.2015.06.015 [arXiv:1501.06795 [hep-ex]].
21. F. Happacher *et al.* [KLOE-2 Collaboration], “Commissioning of the new calorimeters of the KLOE-2 experiment,” *Acta Phys. Polon. B* **46**, no. 1, 87 (2015) doi:10.5506/APhysPolB.46.87 [arXiv:1501.05442 [physics.ins-det]].

22. T. A. Aaltonen *et al.* [CDF Collaboration], “Constraints on Models of the Higgs Boson with Exotic Spin and Parity using Decays to Bottom-Antibottom Quarks in the Full CDF Data Set,” *Phys. Rev. Lett.* **114**, no. 14, 141802 (2015) doi:10.1103/PhysRevLett.114.141802 [arXiv:1501.04875 [hep-ex]].
23. T. Aaltonen *et al.* [CDF Collaboration], “Search for production of an  $\Upsilon(1S)$  meson in association with a  $W$  or  $Z$  boson using the full 1.96 TeV  $p\bar{p}$  collision data set at CDF,” *Phys. Rev. D* **91**, no. 5, 052011 (2015) doi:10.1103/PhysRevD.91.052011 [arXiv:1412.4827 [hep-ex]].
24. T. A. Aaltonen *et al.* [CDF Collaboration], “Measurement of indirect CP-violating asymmetries in  $D^0 \rightarrow K^+K^-$  and  $D^0 \rightarrow \pi^+\pi^-$  decays at CDF,” *Phys. Rev. D* **90**, no. 11, 111103 (2014) doi:10.1103/PhysRevD.90.111103 [arXiv:1410.5435 [hep-ex]].
25. T. A. Aaltonen *et al.* [CDF Collaboration], “Updated Measurement of the Single Top Quark Production Cross Section and  $Vtb$  in the Missing Transverse Energy Plus Jets Topology in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96$  TeV,” *Phys. Rev. D* **93**, no. 3, 032011 (2016) doi:10.1103/PhysRevD.93.032011 [arXiv:1410.4909 [hep-ex]].
26. L. Bartoszek *et al.* [Mu2e Collaboration], “Mu2e Technical Design Report,” arXiv:1501.05241 [physics.ins-det].
27. T. A. Aaltonen *et al.* [CDF Collaboration], “Measurement of the Top-Quark Mass in the All-Hadronic Channel using the full CDF data set,” *Phys. Rev. D* **90**, no. 9, 091101 (2014) doi:10.1103/PhysRevD.90.091101 [arXiv:1409.4906 [hep-ex]].
28. D. Babusci *et al.* [KLOE-2 Collaboration], “Study of the Dalitz decay  $\phi \rightarrow \eta e^+e^-$  with the KLOE detector,” *Phys. Lett. B* **742**, 1 (2015) doi:10.1016/j.physletb.2015.01.011 [arXiv:1409.4582 [hep-ex]].
29. T. A. Aaltonen *et al.* [CDF Collaboration], “Measurement of differential production cross section for  $Z/\gamma^*$  bosons in association with jets in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV,” arXiv:1409.4359 [hep-ex].
30. T. A. Aaltonen *et al.* [CDF Collaboration], “Measurement of the Single Top Quark Production Cross Section and  $-Vtb-$  in Events with One Charged Lepton, Large Missing Transverse Energy, and Jets at CDF,” arXiv:1407.4031 [hep-ex].
31. T. A. Aaltonen *et al.* [CDF Collaboration], “Studies of high-transverse momentum jet substructure and top quarks produced in 1.96 TeV proton-antiproton collisions,” arXiv:1407.3484 [hep-ex].
32. D. Babusci *et al.* [KLOE-2 Collaboration], “Measurement of the absolute branching ratio of the  $K^+ \rightarrow \pi^+\pi^-\pi^+(\gamma)$  decay with the KLOE detector,” [arXiv:1407.2028 [hep-ex]].

33. D. Babusci *et al.* [KLOE-2 Collaboration], “Search for light vector boson production in  $e^+e^- \rightarrow \mu^+\mu^-\gamma$  interactions with the KLOE experiment,” *Phys. Lett. B* **736**, 459 (2014) [arXiv:1404.7772 [hep-ex]].
34. T. A. Aaltonen *et al.* [CDF Collaboration], “Measurement of the inclusive leptonic asymmetry in top-quark pairs that decay to two charged leptons at CDF,” *Phys. Rev. Lett.* **113**, 042001 (2014) [arXiv:1404.3698 [hep-ex]].
35. T. A. Aaltonen *et al.* [CDF Collaboration], “Measurement of  $B(t \rightarrow Wb)/B(t \rightarrow Wq)$  in Top-Quark-Pair Decays Using Dilepton Events and the Full CDF Run II Data Set,” *Phys. Rev. Lett.* **112**, 221801 (2014) [arXiv:1404.3392 [hep-ex]].
36. T. A. Aaltonen *et al.* [CDF Collaboration], “Mass and lifetime measurements of bottom and charm baryons in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV,” *Phys. Rev. D* **89**, 072014 (2014) [arXiv:1403.8126 [hep-ex]].
37. T. A. Aaltonen *et al.* [CDF Collaboration], “Measurements of Direct CP-Violating Asymmetries in Charmless Decays of Bottom Baryons,” arXiv:1403.5586 [hep-ex].
38. G. Pezzullo, J. Budagov, R. Carosi, F. Cervelli, C. Cheng, M. Cordelli, G. Corradi and Y. Davydov *et al.*, “The LYSO crystal calorimeter for the Mu2e experiment,” *JINST* **9**, C03018 (2014).
39. T. A. Aaltonen *et al.* [CDF Collaboration], “Measurement of the ZZ production cross section using the full CDF II data set,” *Phys. Rev. D* **89**, 112001 (2014) [arXiv:1403.2300 [hep-ex]].
40. T. Aaltonen *et al.* [CDF Collaboration], “Invariant-mass distribution of jet pairs produced in association with a  $W$  boson in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV using the full CDF Run II data set,” *Phys. Rev. D* **89**, 092001 (2014) [arXiv:1402.7044 [hep-ex]].
41. T. A. Aaltonen *et al.* [CDF Collaboration], “Study of Top-Quark Production and Decays involving a Tau Lepton at CDF and Limits on a Charged-Higgs Boson Contribution,” *Phys. Rev. D* **89**, 091101 (2014) [arXiv:1402.6728 [hep-ex]].
42. T. A. Aaltonen *et al.* [CDF and D0 Collaborations], “Observation of s-channel production of single top quarks at the Tevatron,” *Phys. Rev. Lett.* **112**, 231803 (2014) [arXiv:1402.5126 [hep-ex]].
43. T. A. Aaltonen *et al.* [CDF Collaboration], “Search for s-channel Single Top Quark Production in the Missing Energy Plus Jets Sample using the Full CDF II Data Set,” *Phys. Rev. Lett.* **112**, 231805 (2014) [arXiv:1402.3756 [hep-ex]].
44. T. A. Aaltonen *et al.* [CDF Collaboration], “Indirect measurement of  $\sin^2 \theta_W$  (or  $M_W$ ) using  $\mu^+\mu^-$  pairs from  $\gamma^*/Z$  bosons produced in  $p\bar{p}$  collisions at a

- center-of-momentum energy of 1.96 TeV,” Phys. Rev. D **89**, 072005 (2014) [arXiv:1402.2239 [hep-ex]].
45. T. A. Aaltonen *et al.* [CDF Collaboration], “Evidence for  $s$ -channel Single-Top-Quark Production in Events with one Charged Lepton and two Jets at CDF,” Phys. Rev. Lett. **112**, 231804 (2014) [arXiv:1402.0484 [hep-ex]].
  46. D. Babusci *et al.* [KLOE-2 Collaboration], “Test of CPT and Lorentz symmetry in entangled neutral kaons with the KLOE experiment,” Phys. Lett. B **730**, 89 (2014) [arXiv:1312.6818 [hep-ex]].
  47. T. A. Aaltonen *et al.* [CDF Collaboration], “First Search for Exotic Z Boson Decays into Photons and Neutral Pions in Hadron Collisions,” Phys. Rev. Lett. **112**, 111803 (2014) [arXiv:1311.3282 [hep-ex]].
  48. T. A. Aaltonen *et al.* [CDF Collaboration], “A precise measurement of the  $W$ -boson mass with the Collider Detector at Fermilab,” Phys. Rev. D **89**, 072003 (2014) [arXiv:1311.0894 [hep-ex]].
  49. T. Aaltonen *et al.* [CDF Collaboration], “Search for a dijet resonance in events with jets and missing transverse energy in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV,” Phys. Rev. D **88**, no. 9, 092004 (2013) [arXiv:1310.7267 [hep-ex]].
  50. E. Czerwinski *et al.* [KLOE/KLOE-2 Collaboration], “KLOE results in flavour physics and prospects for KLOE-2,” Nucl. Phys. Proc. Suppl. **241-242**, 24 (2013).
  51. T. A. Aaltonen *et al.* [CDF Collaboration], “Search for the production of ZW and ZZ boson pairs decaying into charged leptons and jets in  $p\bar{p}$  collisions at  $\sqrt{s}=1.96$  TeV,” Phys. Rev. D **88**, no. 9, 092002 (2013) [arXiv:1310.0086 [hep-ex]].
  52. T. A. Aaltonen *et al.* [CDF and D0 Collaborations], “Combination of measurements of the top-quark pair production cross section from the Tevatron Collider,” Phys. Rev. D **89**, 072001 (2014) [arXiv:1309.7570 [hep-ex]].
  53. T. A. Aaltonen *et al.* [CDF Collaboration], “Search for new physics in trilepton events and limits on the associated chargino-neutralino production at CDF,” Phys. Rev. D **90**, 012011 (2014) [arXiv:1309.7509 [hep-ex]].
  54. T. A. Aaltonen *et al.* [CDF Collaboration], “Study of orbitally excited  $B$  mesons and evidence for a new  $B\pi$  resonance,” Phys. Rev. D **90**, 012013 (2014) [arXiv:1309.5961 [hep-ex]].
  55. T. A. Aaltonen *et al.* [CDF Collaboration], “Observation of  $D^0 - \bar{D}^0$  Mixing Using the CDF II Detector,” Phys. Rev. Lett. **111**, no. 23, 231802 (2013) [arXiv:1309.4078 [hep-ex]].

56. M. Cordelli, E. Dan, S. Giovannella, M. Gatta, F. Happacher, A. Luc, S. Miscetti and A. Saputi *et al.*, “CCALT: A Crystal CALorimeter with Timing for the KLOE-2 upgrade,” Nucl. Instrum. Meth. A **718**, 81 (2013).
57. J. Budagov, R. Carosi, F. Cervelli, C. Cheng, M. Cordelli, Y. Davydov, E. J. Downie and B. Echenard *et al.*, “The calorimeter project for the Mu2e experiment,” Nucl. Instrum. Meth. A **718**, 56 (2013).
58. M. Martemianov *et al.* [KLOE-2 Collaboration], “Status and perspectives of the KLOE-2 experiment,”
59. T. A. Aaltonen *et al.* [CDF Collaboration], “Direct Measurement of the Total Decay Width of the Top Quark,” Phys. Rev. Lett. **111**, no. 20, 202001 (2013) [arXiv:1308.4050 [hep-ex]].
60. T. Aaltonen *et al.* [CDF Collaboration], ‘Production of  $K_S^{0* \pm}(892)$  and  $\phi^0(1020)$  in minimum bias events and  $K_S^0$  and  $\Lambda^0$  in jets in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV” Phys. Rev. D **88**, no. 9, 092005 (2013) [arXiv:1308.3371 [hep-ex]].
61. T. A. Aaltonen *et al.* [CDF Collaboration], “Evidence for a bottom baryon resonance  $\Lambda_b^{*0}$  in CDF data,” Phys. Rev. D **88**, no. 7, 071101 (2013) [arXiv:1308.1760 [hep-ex]].
62. T. A. Aaltonen *et al.* [CDF Collaboration], “Measurement of the leptonic asymmetry in  $t\bar{t}$  events produced in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV,” Phys. Rev. D **88**, no. 7, 072003 (2013) [arXiv:1308.1120 [hep-ex]].
63. T. A. Aaltonen *et al.* [CDF and D0 Collaborations], “Combination of CDF and D0  $W$ -Boson Mass Measurements,” Phys. Rev. D **88**, no. 5, 052018 (2013) [arXiv:1307.7627 [hep-ex]].
64. T. Aaltonen *et al.* [CDF Collaboration], “Indirect measurement of  $\sin^2 \theta_W$  ( $M_W$ ) using  $e^+e^-$  pairs in the Z-boson region with  $p\bar{p}$  collisions at a center-of-momentum energy of 1.96 TeV,” Phys. Rev. D **88**, no. 7, 072002 (2013) [Erratum-ibid. D **88**, no. 7, 079905 (2013)] [arXiv:1307.0770 [hep-ex]].
65. T. Aaltonen *et al.* [CDF Collaboration], “Signature-based search for delayed photons in exclusive photon plus missing transverse energy events from  $p\bar{p}$  collisions with  $\sqrt{s} = 1.96$  TeV,” Phys. Rev. D **88**, no. 3, 031103 (2013) [arXiv:1307.0474 [hep-ex]].
66. P. Moskal *et al.* [KLOE and KLOE-2 Collaborations], “Recent results on hadron physics at KLOE,” arXiv:1306.5740 [hep-ex].
67. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the Differential Cross Section  $d\sigma/d(\cos \theta_t)$  for Top-Quark Pair Production in  $p - \bar{p}$  Collisions at  $\sqrt{s} = 1.96$  TeV,” Phys. Rev. Lett. **111**, no. 18, 182002 (2013) [arXiv:1306.2357 [hep-ex]].

68. T. Aaltonen *et al.* [CDF Collaboration], “Searches for the Higgs boson decaying to  $W^+W^- \rightarrow \ell^+\nu\ell^-\bar{\nu}$  with the CDF II detector,” *Phys. Rev. D* **88**, no. 5, 052012 (2013) [arXiv:1306.0023 [hep-ex]].
69. T. Aaltonen *et al.* [CDF Collaboration], “Top-quark mass measurement in events with jets and missing transverse energy using the full CDF data set,” *Phys. Rev. D* **88**, no. 1, 011101 (2013) [arXiv:1305.3339 [hep-ex]].
70. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the top-quark pair production cross-section in events with two leptons and bottom-quark jets using the full CDF data set,” *Phys. Rev. D* **88**, 091103 (2013) [arXiv:1304.7961 [hep-ex]].
71. T. Aaltonen *et al.* [CDF Collaboration], “Exclusion of exotic top-like quarks with  $-4/3$  electric charge using jet-charge tagging in single-lepton  $t\bar{t}$  events at CDF,” *Phys. Rev. D* **88**, no. 3, 032003 (2013) [arXiv:1304.4141 [hep-ex]].
72. T. Aaltonen *et al.* [CDF and D0 Collaborations], “Higgs Boson Studies at the Tevatron,” *Phys. Rev. D* **88**, no. 5, 052014 (2013) [arXiv:1303.6346 [hep-ex]].
73. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of  $R = \mathcal{B}(t \rightarrow Wb)/\mathcal{B}(t \rightarrow Wq)$  in Top-quark-pair Decays using Lepton+jets Events and the Full CDF Run II Data set,” *Phys. Rev. D* **87**, no. 11, 111101 (2013) [arXiv:1303.6142 [hep-ex]].
74. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the cross section for direct-photon production in association with a heavy quark in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV,” *Phys. Rev. Lett.* **111**, no. 4, 042003 (2013) [arXiv:1303.6136 [hep-ex]].
75. T. Aaltonen *et al.* [CDF Collaboration], “Search for Pair Production of Strongly Interacting Particles Decaying to Pairs of Jets in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96$  TeV,” *Phys. Rev. Lett.* **111**, no. 3, 031802 (2013) [arXiv:1303.2699 [hep-ex]].
76. T. Aaltonen *et al.* [CDF Collaboration], “Search for Supersymmetry with Like-Sign Lepton-Tau Events at CDF,” *Phys. Rev. Lett.* **110**, no. 20, 201802 (2013) [arXiv:1302.4491 [hep-ex]].
77. D. Babusci *et al.* [KLOE Collaboration], “A new limit on the CP violating decay  $K_S \rightarrow 3\pi^0$  with the KLOE experiment,” *Phys. Lett. B* **723**, 54 (2013) [arXiv:1301.7623 [hep-ex]].
78. T. Aaltonen *et al.* [CDF Collaboration], “Search for  $B_s \rightarrow \mu^+\mu^-$  and  $B_d \rightarrow \mu^+\mu^-$  decays with the full CDF Run II data set,” *Phys. Rev. D* **87**, 072003 (2013) [arXiv:1301.7048 [hep-ex]].
79. T. Aaltonen *et al.* [CDF Collaboration], “Combination fo Searches for the Higgs Boson Using the Full CDF Data Set,” *Phys. Rev. D* **88**, no. 5, 052013 (2013) [arXiv:1301.6668 [hep-ex]].

80. T. Aaltonen *et al.* [CDF Collaboration], “Updated search for the standard model Higgs boson in events with jets and missing transverse energy using the full CDF data set,” *Phys. Rev. D* **87**, 052008 (2013) [arXiv:1301.4440 [hep-ex]].
81. D. Babusci *et al.* [KLOE Collaboration], “Precision measurement of  $\sigma(e^+e^- \rightarrow \pi^+\pi^-\gamma)/\sigma(e^+e^- \rightarrow \mu^+\mu^-\gamma)$  and determination of the  $\pi^+\pi^-$  contribution to the muon anomaly with the KLOE detector,” *Phys. Lett. B* **720**, 336 (2013) [arXiv:1212.4524 [hep-ex]].
82. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the cross section for prompt isolated diphoton production using the full CDF Run II data sample,” *Phys. Rev. Lett.* **110**, no. 10, 101801 (2013) [arXiv:1212.4204 [hep-ex]].
83. T. Aaltonen *et al.* [CDF Collaboration], “Search for a two-Higgs-boson doublet using a simplified model in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV,” *Phys. Rev. Lett.* **110**, no. 12, 121801 (2013) [arXiv:1212.3837 [hep-ex]].
84. R. J. Abrams *et al.* [Mu2e Collaboration], “Mu2e Conceptual Design Report,” arXiv:1211.7019 [physics.ins-det].
85. T. Aaltonen *et al.* [CDF Collaboration], “Search for Resonant Top-Antitop Production in the Lepton Plus Jets Decay Mode Using the Full CDF Data Set,” *Phys. Rev. Lett.* **110**, no. 12, 121802 (2013) [arXiv:1211.5363 [hep-ex]].
86. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of  $W$ -Boson Polarization in Top-quark Decay using the Full CDF Run II Data Set,” *Phys. Rev. D* **87**, 031104 (2013) [arXiv:1211.4523 [hep-ex]].
87. D. Babusci *et al.* [KLOE-2 Collaboration], “Measurement of  $\eta$  meson production in  $\gamma\gamma$  interactions and  $\Gamma(\eta \rightarrow \gamma\gamma)$  with the KLOE detector,” *JHEP* **1301**, 119 (2013) [*JHEP* **1301**, 119 (2013)] [arXiv:1211.1845 [hep-ex]].
88. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the top quark forward-backward production asymmetry and its dependence on event kinematic properties,” *Phys. Rev. D* **87**, 092002 (2013) [arXiv:1211.1003 [hep-ex]].
89. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the mass difference between top and antitop quarks,” *Phys. Rev. D* **87**, no. 5, 052013 (2013) [arXiv:1210.6131 [hep-ex]].
90. T. Aaltonen *et al.* [CDF Collaboration], “Search for a heavy vector boson decaying to two gluons in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV,” *Phys. Rev. D* **86**, 112002 (2012) [arXiv:1210.5686 [hep-ex]].
91. D. Babusci *et al.* [KLOE-2 Collaboration], “Limit on the production of a light vector gauge boson in phi meson decays with the KLOE detector,” *Phys. Lett. B* **720**, 111 (2013) [arXiv:1210.3927 [hep-ex]].

92. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the  $B_c^-$  meson lifetime in the decay  $B_c^- \rightarrow J/\psi \pi^-$ ,” Phys. Rev. D **87**, 011101 (2013) [arXiv:1210.2366 [hep-ex]].
93. D. Babusci *et al.* [KLOE Collaboration], “Measurement of  $\Gamma(\eta \rightarrow \pi^+ \pi^- \gamma)/\Gamma(\eta \rightarrow \pi^+ \pi^- \pi^0)$  with the KLOE Detector,” Phys. Lett. B **718**, 910 (2013) [arXiv:1209.4611 [hep-ex]].
94. T. Aaltonen *et al.* [CDF Collaboration], “Observation of the Production of a W Boson in Association with a Single Charm Quark,” Phys. Rev. Lett. **110**, no. 7, 071801 (2013) [arXiv:1209.1921 [hep-ex]].
95. T. Aaltonen *et al.* [CDF Collaboration], “Measurements of the Top-quark Mass and the  $t\bar{t}$  Cross Section in the Hadronic  $\tau+$  Jets Decay Channel at  $\sqrt{s} = 1.96$  TeV,” Phys. Rev. Lett. **109**, 192001 (2012) [arXiv:1208.5720 [hep-ex]].
96. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the Bottom-Strange Meson Mixing Phase in the Full CDF Data Set,” Phys. Rev. Lett. **109**, 171802 (2012) [arXiv:1208.2967 [hep-ex]].
97. T. Aaltonen *et al.* [CDF Collaboration], “Search for the standard model Higgs boson produced in association with top quarks using the full CDF data set,” Phys. Rev. Lett. **109**, 181802 (2012) [arXiv:1208.2662 [hep-ex]].
98. T. Aaltonen *et al.* [CDF Collaboration], “Search for the Higgs boson in the all-hadronic final state using the full CDF data set,” JHEP **1302**, 004 (2013) [arXiv:1208.6445 [hep-ex]].
99. T. Aaltonen *et al.* [CDF Collaboration], “Transverse momentum cross section of  $e^+e^-$  pairs in the  $Z$ -boson region from  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV,” Phys. Rev. D **86**, 052010 (2012) [arXiv:1207.7138].
100. T. Aaltonen *et al.* [CDF Collaboration], “Precision Top-Quark Mass Measurements at CDF,” Phys. Rev. Lett. **109**, 152003 (2012) [arXiv:1207.6758 [hep-ex]].
101. T. Aaltonen *et al.* [CDF and D0 Collaborations], “Evidence for a particle produced in association with weak bosons and decaying to a bottom-antibottom quark pair in Higgs boson searches at the Tevatron,” Phys. Rev. Lett. **109**, 071804 (2012) [arXiv:1207.6436 [hep-ex]].
102. T. Aaltonen *et al.* [CDF Collaboration], “Search for a Higgs boson in the diphoton final state using the full CDF data set from proton-antiproton collisions at  $\sqrt{s} = 1.96$  TeV,” Phys. Lett. B **717**, 173 (2012) [arXiv:1207.6386 [hep-ex]].
103. T. Aaltonen *et al.* [CDF Collaboration], “Novel inclusive search for the Higgs boson in the four-lepton final state at CDF,” Phys. Rev. D **86**, 072012 (2012) [arXiv:1207.5016 [hep-ex]].

104. T. Aaltonen *et al.* [CDF and D0 Collaborations], “Search for Neutral Higgs Bosons in Events with Multiple Bottom Quarks at the Tevatron,” *Phys. Rev. D* **86**, 091101 (2012) [arXiv:1207.2757 [hep-ex]].
105. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the difference of CP-violating asymmetries in  $D^0 \rightarrow K^+K^-$  and  $D^0 \rightarrow \pi^+\pi^-$  decays at CDF,” *Phys. Rev. Lett.* **109**, 111801 (2012) [arXiv:1207.2158 [hep-ex]].
106. T. Aaltonen *et al.* [CDF Collaboration], “Search for the standard model Higgs boson decaying to a  $b\bar{b}$  pair in events with no charged leptons and large missing transverse energy using the full CDF data set,” *Phys. Rev. Lett.* **109**, 111805 (2012) [arXiv:1207.1711 [hep-ex]].
107. T. Aaltonen *et al.* [CDF Collaboration], “Combined search for the standard model Higgs boson decaying to a  $b\bar{b}$  pair using the full CDF data set,” *Phys. Rev. Lett.* **109**, 111802 (2012) [arXiv:1207.1707 [hep-ex]].
108. T. Aaltonen *et al.* [CDF Collaboration], “Search for the standard model Higgs boson decaying to a  $b\bar{b}$  pair in events with two oppositely-charged leptons using the full CDF data set,” *Phys. Rev. Lett.* **109**, 111803 (2012) [arXiv:1207.1704 [hep-ex]].
109. T. Aaltonen *et al.* [CDF Collaboration], “Search for the standard model Higgs boson decaying to a  $b\bar{b}$  pair in events with one charged lepton and large missing transverse energy using the full CDF data set,” *Phys. Rev. Lett.* **109**, 111804 (2012) [arXiv:1207.1703 [hep-ex]].
110. T. Aaltonen *et al.* [CDF and D0 Collaborations], “Combination of the top-quark mass measurements from the Tevatron collider,” *Phys. Rev. D* **86**, 092003 (2012) [arXiv:1207.1069 [hep-ex]].
111. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of CP-violation asymmetries in  $D^0 \rightarrow K_S\pi^+\pi^-$ ,” *Phys. Rev. D* **86**, 032007 (2012) [arXiv:1207.0825 [hep-ex]].
112. T. Aaltonen *et al.* [CDF Collaboration], “Search for the standard model Higgs boson produced in association with a  $W^\pm$  boson with  $7.5 \text{ fb}^{-1}$  integrated luminosity at CDF,” *Phys. Rev. D* **86**, 032011 (2012) [arXiv:1206.5063 [hep-ex]].
113. T. Aaltonen *et al.* [CDF Collaboration], “Diffractive Dijet Production in  $\bar{p}p$  Collisions at  $\sqrt{s} = 1.96 \text{ TeV}$ ,” *Phys. Rev. D* **86**, 032009 (2012) [arXiv:1206.3955 [hep-ex]].
114. T. Aaltonen *et al.* [CDF Collaboration], “ $W$  boson polarization measurement in the  $t\bar{t}$  dilepton channel using the CDF II Detector,” *Phys. Lett. B* **722**, 48 (2013) [arXiv:1205.0354 [hep-ex]].

115. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of  $B_s^0 \rightarrow D_s^{(*)+} D_s^{(*)-}$  Branching Ratios,” Phys. Rev. Lett. **108**, 201801 (2012) [arXiv:1204.0536 [hep-ex]].
116. T. Aaltonen *et al.* [CDF Collaboration], “Search for the standard model Higgs boson produced in association with a  $Z$  Boson in 7.9 fb<sup>-1</sup> of p anti-p collisions at  $\sqrt{s} = 1.96$  TeV,” Phys. Lett. B **715**, 98 (2012) [arXiv:1203.5815 [hep-ex]].
117. T. Aaltonen *et al.* [CDF Collaboration], “Search for Scalar Top Quark Production in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96$  TeV,” JHEP **1210**, 158 (2012) [arXiv:1203.4171 [hep-ex]].
118. T. Aaltonen *et al.* [CDF Collaboration], “Search for a heavy particle decaying to a top quark and a light quark in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV,” Phys. Rev. Lett. **108**, 211805 (2012) [arXiv:1203.3894 [hep-ex]].
119. T. Aaltonen *et al.* [CDF Collaboration], “A Search for dark matter in events with one jet and missing transverse energy in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV,” Phys. Rev. Lett. **108**, 211804 (2012) [arXiv:1203.0742 [hep-ex]].
120. T. Aaltonen *et al.* [CDF Collaboration], “Precise measurement of the  $W$ -boson mass with the CDF II detector,” Phys. Rev. Lett. **108**, 151803 (2012) [arXiv:1203.0275 [hep-ex]].
121. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the  $WZ$  Cross Section and Triple Gauge Couplings in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96$  TeV,” Phys. Rev. D **86**, 031104 (2012) [arXiv:1202.6629 [hep-ex]].
122. T. Aaltonen *et al.* [CDF Collaboration], “Search for a dark matter candidate produced in association with a single top quark in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV,” Phys. Rev. Lett. **108**, 201802 (2012) [arXiv:1202.5653 [hep-ex]].
123. T. Aaltonen *et al.* [CDF and D0 Collaborations], “Combination of CDF and D0 measurements of the  $W$  boson helicity in top quark decays,” Phys. Rev. D **85**, 071106 (2012) [arXiv:1202.5272 [hep-ex]].
124. T. Aaltonen *et al.* [CDF Collaboration], “Search for anomalous production of multiple leptons in association with  $W$  and  $Z$  bosons at CDF,” Phys. Rev. D **85**, 092001 (2012) [arXiv:1202.1260 [hep-ex]].
125. T. Aaltonen *et al.* [CDF Collaboration], “Search for a low mass Standard Model Higgs boson in the  $\tau - \tau$  decay channel in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV,” Phys. Rev. Lett. **108**, 181804 (2012) [arXiv:1201.4880 [hep-ex]].
126. M. Cordelli, S. Giovannella, F. Happacher, A. Luca’, S. Miscetti, A. Saputi, I. Sarra and G. Pileggi *et al.*, “Test and Simulation of a LYSO+APD matrix with a tagged Photon Beam from 40-MeV to 300-MeV,” J. Phys. Conf. Ser. **404**, 012027 (2012).

127. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the Top Quark Mass in the All-Hadronic Mode at CDF,” *Phys. Lett. B* **714**, 24 (2012) [arXiv:1112.4891 [hep-ex]].
128. T. Aaltonen *et al.* [CDF Collaboration], “Search for Standard Model Higgs Boson Production in Association with a  $W$  Boson Using a Matrix Element Technique at CDF in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96$  TeV,” *Phys. Rev. D* **85**, 072001 (2012) [arXiv:1112.4358 [hep-ex]].
129. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the branching fraction  $\mathcal{B}(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^- \pi^+ \pi^-)$  at CDF,” *Phys. Rev. D* **85**, 032003 (2012) [arXiv:1112.3334 [hep-ex]].
130. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the masses and widths of the bottom baryons  $\Sigma_b^{+-}$  and  $\Sigma_b^{*+-}$ ,” *Phys. Rev. D* **85**, 092011 (2012) [arXiv:1112.2808 [hep-ex]].
131. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of  $ZZ$  production in leptonic final states at  $\sqrt{s}$  of 1.96 TeV at CDF,” *Phys. Rev. Lett.* **108**, 101801 (2012) [arXiv:1112.2978 [hep-ex]].
132. T. Aaltonen *et al.* [CDF Collaboration], “Search for Standard Model Higgs Boson Production in Association with a  $W$  Boson at CDF,” *Phys. Rev. D* **85**, 052002 (2012) [arXiv:1112.1930 [hep-ex]].
133. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the CP-Violating Phase  $\beta_s^{J/\Psi\phi}$  in  $B_s^0 \rightarrow J/\Psi\phi$  Decays with the CDF II Detector,” *Phys. Rev. D* **85**, 072002 (2012) [arXiv:1112.1726 [hep-ex]].
134. T. Aaltonen *et al.* [CDF Collaboration], “Measurements of Angular Distributions of Muons From  $\Upsilon$  Meson Decays in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96$  TeV,” *Phys. Rev. Lett.* **108**, 151802 (2012) [arXiv:1112.1591 [hep-ex]].
135. T. Aaltonen *et al.* [CDF Collaboration], “Search for new phenomena in events with two  $Z$  bosons and missing transverse momentum in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV,” *Phys. Rev. D* **85**, 011104 (2012) [arXiv:1112.1577 [hep-ex]].
136. T. Aaltonen *et al.* [CDF Collaboration], “Observation of Exclusive Gamma Gamma Production in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96$  TeV,” *Phys. Rev. Lett.* **108**, 081801 (2012) [arXiv:1112.0858 [hep-ex]].
137. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of CP-violating asymmetries in  $D^0 \rightarrow \pi^+\pi^-$  and  $D^0 \rightarrow K^+K^-$  decays at CDF,” *Phys. Rev. D* **85**, 012009 (2012) [arXiv:1111.5023 [hep-ex]].
138. T. Aaltonen *et al.* [CDF Collaboration], “An additional study of multi-muon events produced in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV,” *Phys. Lett. B* **710**, 278 (2012) [arXiv:1111.5242 [hep-ex]].

139. T. Aaltonen *et al.* [CDF Collaboration], “Search for high-mass resonances decaying into  $ZZ$  in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV,” *Phys. Rev. D* **85**, 012008 (2012) [arXiv:1111.3432 [hep-ex]].
140. T. Aaltonen *et al.* [CDF Collaboration], “Evidence for the charmless annihilation decay mode  $B_s^0 \rightarrow \pi^+\pi^-$ ,” *Phys. Rev. Lett.* **108**, 211803 (2012) [arXiv:1111.0485 [hep-ex]].
141. F. Archilli, D. Babusci, D. Badoni, I. Balwierz, G. Bencivenni, C. Bini, C. Bloise and V. Bocci *et al.*, “Search for a vector gauge boson in phi meson decays with the KLOE detector,” *Phys. Lett. B* **706**, 251 (2012) [arXiv:1110.0411 [hep-ex]].
142. T. Aaltonen *et al.* [CDF Collaboration], “Search for a Higgs Boson in the Diphoton Final State in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96$  TeV,” *Phys. Rev. Lett.* **108**, 011801 (2012) [arXiv:1109.4427 [hep-ex]].
143. T. Aaltonen *et al.* [CDF Collaboration], “Search for heavy metastable particles decaying to jet pairs in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV,” *Phys. Rev. D* **85**, 012007 (2012) [arXiv:1109.3136 [hep-ex]].
144. T. Aaltonen *et al.* [CDF Collaboration], “Top-quark mass measurement using events with missing transverse energy and jets at CDF,” *Phys. Rev. Lett.* **107**, 232002 (2011) [arXiv:1109.1490 [hep-ex]].
145. T. Aaltonen *et al.* [CDF Collaboration], “Measurements of branching fraction ratios and CP-asymmetries in suppressed  $B^- \rightarrow D(\rightarrow K^+\pi^-)K^-$  and  $B^- \rightarrow D(\rightarrow K^+\pi^-)\pi^-$  decays,” *Phys. Rev. D* **84**, 091504 (2011) [arXiv:1108.5765 [hep-ex]].
146. T. Aaltonen *et al.* [CDF Collaboration], “Search for resonant production of  $t\bar{t}$  decaying to jets in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV,” *Phys. Rev. D* **84**, 072003 (2011) [arXiv:1108.4755 [hep-ex]].
147. T. Aaltonen *et al.* [CDF Collaboration], “Search for  $WZ + ZZ$  production with MET + jets with  $b$  enhancement at  $\sqrt{s} = 1.96$  TeV,” *Phys. Rev. D* **85**, 012002 (2012) [arXiv:1108.2060 [hep-ex]].
148. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the top-quark mass in the lepton+jets channel using a matrix element technique with the CDF II detector,” *Phys. Rev. D* **84**, 071105 (2011) [arXiv:1108.1601 [hep-ex]].
149. T. Aaltonen *et al.* [CDF Collaboration], “Measurements of the Angular Distributions in the Decays  $B \rightarrow K^{(*)}\mu^+\mu^-$  at CDF,” *Phys. Rev. Lett.* **108**, 081807 (2012) [arXiv:1108.0695 [hep-ex]].
150. T. Aaltonen *et al.* [CDF Collaboration], “Search for new physics in high  $p_T$  like-sign dilepton events at CDF II,” *Phys. Rev. Lett.* **107**, 181801 (2011) [arXiv:1108.0101 [hep-ex]].

151. F. Ambrosino *et al.* [KLOE and 2 Collaborations], “Measurement of  $\Gamma(\eta \rightarrow \pi^+\pi^-\gamma)/\Gamma(\eta \rightarrow \pi^+\pi^-\pi^0)$  with KLOE experiment,” arXiv:1107.5733 [hep-ex].
152. T. Aaltonen *et al.* [CDF Collaboration], “A Search for resonant production of  $t\bar{t}$  pairs in  $4.8 \text{ fb}^{-1}$  of integrated luminosity of  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96 \text{ TeV}$ ,” Phys. Rev. D **84**, 072004 (2011) [arXiv:1107.5063 [hep-ex]].
153. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of Polarization and Search for CP-Violation in  $B_s^0 \rightarrow \phi\phi$  Decays,” Phys. Rev. Lett. **107**, 261802 (2011) [arXiv:1107.4999 [hep-ex]].
154. T. Aaltonen *et al.* [CDF Collaboration], “Search for a Heavy Top-Like Quark in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96 \text{ TeV}$ ,” Phys. Rev. Lett. **107**, 261801 (2011) [arXiv:1107.3875 [hep-ex]].
155. T. Aaltonen *et al.* [CDF Collaboration], “Observation of the  $\Xi_b^0$  Baryon,” Phys. Rev. Lett. **107**, 102001 (2011) [arXiv:1107.4015 [hep-ex]].
156. T. Aaltonen *et al.* [CDF Collaboration], “Search for New  $T'$  Particles in Final States with Large Jet Multiplicities and Missing Transverse Energy in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96 \text{ TeV}$ ,” Phys. Rev. Lett. **107**, 191803 (2011) [arXiv:1107.3574 [hep-ex]].
157. T. Aaltonen *et al.* [CDF Collaboration], “Observation of the Baryonic Flavor-Changing Neutral Current Decay  $\Lambda_b \rightarrow \Lambda\mu^+\mu^-$ ,” Phys. Rev. Lett. **107**, 201802 (2011) [arXiv:1107.3753 [hep-ex]].
158. F. Archilli *et al.* [KLOE-2 Collaboration], “ $\gamma\gamma$  physics with the KLOE experiment,” arXiv:1107.3782 [hep-ex].
159. T. Aaltonen *et al.* [CDF Collaboration], “Search for  $B_s^{0+-}$  and  $B_0^{+-}$  Decays with CDF II,” Phys. Rev. Lett. **107**, 191801 (2011) [Publisher-note **107**, 239903 (2011)] [arXiv:1107.2304 [hep-ex]].
160. T. Aaltonen *et al.* [CDF Collaboration], “Study of Substructure of High Transverse Momentum Jets Produced in Proton-Antiproton Collisions at  $\sqrt{s} = 1.96 \text{ TeV}$ ,” Phys. Rev. D **85**, 091101 (2012) [arXiv:1106.5952 [hep-ex]].
161. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the Cross Section for Prompt Isolated Diphoton Production in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96 \text{ TeV}$ ,” Phys. Rev. Lett. **107**, 102003 (2011) [arXiv:1106.5123 [hep-ex]].
162. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the Cross Section for Prompt Isolated Diphoton Production in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96 \text{ TeV}$ ,” Phys. Rev. D **84**, 052006 (2011) [arXiv:1106.5131 [hep-ex]].
163. T. Aaltonen *et al.* [CDF Collaboration], “Search for Higgs Bosons Produced in Association with  $b$ -quarks,” Phys. Rev. D **85**, 032005 (2012) [arXiv:1106.4782 [hep-ex]].

164. T. Aaltonen *et al.* [CDF Collaboration], “Evidence for  $t\bar{t}\gamma$  Production and Measurement of  $\sigma_{t\bar{t}\gamma}/\sigma_{t\bar{t}}$ ,” Phys. Rev. D **84**, 031104 (2011) [arXiv:1106.3970 [hep-ex]].
165. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of branching ratio and  $B_s^0$  lifetime in the decay  $B_s^0 \rightarrow J/\psi f_0(980)$  at CDF,” Phys. Rev. D **84**, 052012 (2011) [arXiv:1106.3682 [hep-ex]].
166. M. Cordelli, F. Happacher, M. Martini, S. Miscetti, I. Sarra, M. Schioppa and S. Stucci, “CCALT: A crystal calorimeter for the KLOE-2 experiment,” J. Phys. Conf. Ser. **293**, 012010 (2011).
167. F. Ambrosino *et al.* [KLOE-2 Collaboration], “Observation of the rare  $\eta \rightarrow e^+e^-e^+e^-$  decay with the KLOE experiment,” Phys. Lett. B **702**, 324 (2011) [arXiv:1105.6067 [hep-ex]].
168. T. Aaltonen *et al.* [CDF Collaboration], “Measurements of the properties of  $\Lambda_c(2595)$ ,  $\Lambda_c(2625)$ ,  $\Sigma_c(2455)$ , and  $\Sigma_c(2520)$  baryons,” Phys. Rev. D **84**, 012003 (2011) [arXiv:1105.5995 [hep-ex]].
169. T. Aaltonen *et al.* [CDF Collaboration], “Improved determination of the sample composition of dimuon events produced in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV,” Eur. Phys. J. C **71**, 1720 (2011) [arXiv:1105.5787 [hep-ex]].
170. T. Aaltonen *et al.* [CDF Collaboration], “First Search for Multijet Resonances in  $\sqrt{s} = 1.96$  TeV  $p\bar{p}$  Collisions,” Phys. Rev. Lett. **107**, 042001 (2011) [arXiv:1105.2815 [hep-ex]].
171. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the  $t\bar{t}$  production cross section in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV using events with large Missing  $E_T$  and jets,” Phys. Rev. D **84**, 032003 (2011) [arXiv:1105.1806 [hep-ex]].
172. T. Aaltonen *et al.* [CDF Collaboration], “Top quark mass measurement using the template method at CDF,” Phys. Rev. D **83**, 111101 (2011) [arXiv:1105.0192 [hep-ex]].
173. T. Aaltonen *et al.* [CDF Collaboration], “Search for the Rare Radiative Decay:  $W \rightarrow \pi\gamma$  in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96$  TeV,” Phys. Rev. D **85**, 032001 (2012) [arXiv:1104.1585 [hep-ex]].
174. T. Aaltonen *et al.* [CDF Collaboration], “Invariant Mass Distribution of Jet Pairs Produced in Association with a  $W$  boson in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96$  TeV,” Phys. Rev. Lett. **106**, 171801 (2011) [arXiv:1104.0699 [hep-ex]].
175. T. Aaltonen *et al.* [CDF Collaboration], “Search for a Very Light CP-Odd Higgs Boson in Top Quark Decays from  $p^-p$  Collisions at 1.96 TeV,” Phys. Rev. Lett. **107**, 031801 (2011) [arXiv:1104.5701 [hep-ex]].

176. T. Aaltonen *et al.* [CDF Collaboration], “Measurements of Direct CP Violating Asymmetries in Charmless Decays of Strange Bottom Mesons and Bottom Baryons,” *Phys. Rev. Lett.* **106**, 181802 (2011) [arXiv:1103.5762 [hep-ex]].
177. T. Aaltonen *et al.* [CDF Collaboration], “First Measurement of the Angular Coefficients of Drell-Yan  $e^+e^-$  pairs in the Z Mass Region from  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96$  TeV,” *Phys. Rev. Lett.* **106**, 241801 (2011) [arXiv:1103.5699 [hep-ex]].
178. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of Event Shapes in Proton-Antiproton Collisions at Center-of-Mass Energy 1.96 TeV,” *Phys. Rev. D* **83**, 112007 (2011) [arXiv:1103.5143 [hep-ex]].
179. T. Aaltonen *et al.* [CDF Collaboration], “Search for New Dielectron Resonances and Randall-Sundrum Gravitons at the Collider Detector at Fermilab,” *Phys. Rev. Lett.* **107**, 051801 (2011) [arXiv:1103.4650 [hep-ex]].
180. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the Top Pair Production Cross Section in the Lepton + Jets Channel Using a Jet Flavor Discriminant,” *Phys. Rev. D* **84**, 031101 (2011) [arXiv:1103.4821 [hep-ex]].
181. T. Aaltonen *et al.* [CDF and D0 Collaborations], “Combined CDF and D0 Upper Limits on Standard Model Higgs Boson Production with up to  $8.2 \text{ fb}^{-1}$  of Data,” arXiv:1103.3233 [hep-ex].
182. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the mass difference between  $t$  and  $\bar{t}$  quarks,” *Phys. Rev. Lett.* **106**, 152001 (2011) [arXiv:1103.2782 [hep-ex]].
183. T. Aaltonen *et al.* [CDF Collaboration], “Limits on Anomalous Trilinear Gauge Couplings in  $Z\gamma$  Events from  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96$  TeV,” *Phys. Rev. Lett.* **107**, 051802 (2011) [arXiv:1103.2990 [hep-ex]].
184. T. Aaltonen *et al.* [CDF Collaboration], “Search for Production of Heavy Particles Decaying to Top Quarks and Invisible Particles in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV,” *Phys. Rev. Lett.* **106**, 191801 (2011) [arXiv:1103.2482 [hep-ex]].
185. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the  $B_s$  Lifetime in Fully and Partially Reconstructed  $B_s \rightarrow D_s^-(\phi\pi^-)X$  Decays in  $\bar{p} - p$  Collisions at  $\sqrt{s} = 1.96$  TeV,” *Phys. Rev. Lett.* **107**, 272001 (2011) [arXiv:1103.1864 [hep-ex]].
186. T. Aaltonen *et al.* [CDF Collaboration], “Search for New Heavy Particles Decaying to  $ZZ \rightarrow \ell\ell\ell, \ell\ell jj$  in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96$  TeV,” *Phys. Rev. D* **83**, 112008 (2011) [arXiv:1102.4566 [hep-ex]].
187. T. Aaltonen *et al.* [CDF Collaboration], “Observation of  $B_s^0 \rightarrow J/\psi K^{*0}(892)$  and  $B_s^0 \rightarrow J/\psi K_S^0$  Decays,” *Phys. Rev. D* **83**, 052012 (2011) [arXiv:1102.1961 [hep-ex]].

188. T. Aaltonen *et al.* [CDF Collaboration], “Search for the Higgs boson in the all-hadronic final state using the CDF II detector,” Phys. Rev. D **84**, 052010 (2011) [arXiv:1102.0024 [hep-ex]].
189. T. Aaltonen *et al.* [CDF Collaboration], “Observation of the  $Y(4140)$  structure in the  $J/\psi \phi$  Mass Spectrum in  $B^\pm \rightarrow J/\psi \phi K$  decays,” arXiv:1101.6058 [hep-ex].
190. T. Aaltonen *et al.* [CDF Collaboration], “Search for heavy bottom-like quarks decaying to an electron or muon and jets in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV,” Phys. Rev. Lett. **106**, 141803 (2011) [arXiv:1101.5728 [hep-ex]].
191. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the Top Quark Mass in the Lepton+Jets Channel Using the Lepton Transverse Momentum,” Phys. Lett. B **698**, 371 (2011) [arXiv:1101.4926 [hep-ex]].
192. T. Aaltonen *et al.* [CDF Collaboration], “Search for High Mass Resonances Decaying to Muon Pairs in  $\sqrt{s} = 1.96$  TeV  $p\bar{p}$  Collisions,” Phys. Rev. Lett. **106**, 121801 (2011) [arXiv:1101.4578 [hep-ex]].
193. T. Aaltonen *et al.* [CDF Collaboration], “Production of  $\Lambda$ ,  $\bar{\Lambda}^0$ ,  $\Xi^\pm$  and  $\Omega^\pm$  Hyperons in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96$  TeV,” Phys. Rev. D **86**, 012002 (2012) [arXiv:1101.2996 [hep-ex]].
194. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the Forward-Backward Asymmetry in the  $B \rightarrow K^{(*)} \mu^+ \mu^-$  Decay and First Observation of the  $B_s^0 \rightarrow \phi \mu^+ \mu^-$  Decay,” Phys. Rev. Lett. **106**, 161801 (2011) [arXiv:1101.1028 [hep-ex]].
195. T. Aaltonen *et al.* [CDF Collaboration], “Evidence for a Mass Dependent Forward-Backward Asymmetry in Top Quark Pair Production,” Phys. Rev. D **83**, 112003 (2011) [arXiv:1101.0034 [hep-ex]].
196. T. Aaltonen *et al.* [CDF Collaboration], “Search for a New Heavy Gauge Boson  $W'$  with Electron + missing ET Event Signature in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV,” Phys. Rev. D **83**, 031102 (2011) [arXiv:1012.5145 [hep-ex]].
197. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of  $t\bar{t}$  Spin Correlation in  $p\bar{p}$  Collisions Using the CDF II Detector at the Tevatron,” Phys. Rev. D **83**, 031104 (2011) [arXiv:1012.3093 [hep-ex]].
198. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of  $b$  hadron lifetimes in exclusive decays containing a  $J/\psi$  in  $p^-$  pbar collisions at  $\sqrt{s} = 1.96$  TeV,” Phys. Rev. Lett. **106**, 121804 (2011) [arXiv:1012.3138 [hep-ex]].
199. T. Aaltonen *et al.* [CDF Collaboration], “Search for Randall-Sundrum Gravitons in the Diphoton Channel at CDF,” Phys. Rev. D **83**, 011102 (2011) [arXiv:1012.2795 [hep-ex]].

200. T. Aaltonen *et al.* [CDF Collaboration], “Top Quark Mass Measurement in the Lepton + Jets Channel Using a Matrix Element Method and *in situ* Jet Energy Calibration,” Phys. Rev. Lett. **105**, 252001 (2010) [arXiv:1010.4582 [hep-ex]].
201. T. Aaltonen *et al.* [CDF Collaboration], “Improved Search for a Higgs Boson Produced in Association with  $Z \rightarrow \ell^+\ell^-$  in proton antiproton Collisions at  $\sqrt{s} = 1.96$  TeV,” Phys. Rev. Lett. **105**, 251802 (2010) [arXiv:1009.3047 [hep-ex]].
202. T. Aaltonen *et al.* [CDF Collaboration], “Search for the supersymmetric partner of the top quark in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV,” Phys. Rev. D **82**, 092001 (2010) [arXiv:1009.0266 [hep-ex]].
203. E. Czerwinski *et al.* [on behalf of KLOE-2 Collaboration], “Hadron physics with KLOE-2,” Nucl. Phys. Proc. Suppl. **207-208**, 137 (2010) [arXiv:1009.0113 [nucl-ex]].
204. T. Aaltonen *et al.* [CDF Collaboration], “Updated Search for the Flavor-Changing Neutral-Current Decay  $D^0 \rightarrow \mu^+\mu^-$ ,” Phys. Rev. D **82**, 091105 (2010) [arXiv:1008.5077 [hep-ex]].
205. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the  $WW + WZ$  Production Cross Section Using a Matrix Element Technique in Lepton + Jets Events,” Phys. Rev. D **82**, 112001 (2010) [arXiv:1008.4404 [hep-ex]].
206. T. Aaltonen *et al.* [CDF Collaboration], “Direct Top-Quark Width Measurement CDF,” Phys. Rev. Lett. **105**, 232003 (2010) [arXiv:1008.3891 [hep-ex]].
207. D. Babusci, C. Bini, F. Bossi, G. Isidori, D. Moricciani, F. Nguyen, P. Raimondi and G. Venanzoni *et al.*, “Proposal for taking data with the KLOE-2 detector at the DAΦNE collider upgraded in energy,” arXiv:1007.5219 [hep-ex].
208. T. Aaltonen *et al.* [CDF Collaboration], “Diffractive W and Z Production at the Fermilab Tevatron,” Phys. Rev. D **82**, 112004 (2010) [arXiv:1007.5048 [hep-ex]].
209. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the  $t\bar{t}$  Production Cross Section with an *in situ* Calibration of  $b$ -jet Identification Efficiency,” Phys. Rev. D **83**, 071102 (2011) [arXiv:1007.4423 [hep-ex]].
210. T. Aaltonen *et al.* [CDF Collaboration], “Exclusion of an Exotic Top Quark with  $-4/3$  Electric Charge Using Soft Lepton Tagging,” Phys. Rev. Lett. **105**, 101801 (2010) [arXiv:1006.4597 [hep-ex]].
211. T. Aaltonen *et al.* [CDF Collaboration], “Study of multi-muon events produced in  $p\bar{p}$  interactions at  $\sqrt{s} = 1.96$  TeV,” Eur. Phys. J. C **68**, 109 (2010).
212. T. Aaltonen *et al.* [CDF Collaboration], “Search for the Production of Scalar Bottom Quarks in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV,” Phys. Rev. Lett. **105**, 081802 (2010) [arXiv:1005.3600 [hep-ex]].

213. T. Aaltonen *et al.* [CDF and D0 Collaborations], “Combined Tevatron upper limit on  $gg \rightarrow H \rightarrow W^+W^-$  and constraints on the Higgs boson mass in fourth-generation fermion models,” *Phys. Rev. D* **82**, 011102 (2010) [arXiv:1005.3216 [hep-ex]].
214. T. Aaltonen *et al.* [CDF Collaboration], “Search for  $WW$  and  $WZ$  resonances decaying to electron, missing  $E_T$ , and two jets in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV,” *Phys. Rev. Lett.* **104**, 241801 (2010) [arXiv:1004.4946 [hep-ex]].
215. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the  $B^-$  lifetime using a simulation free approach for trigger bias correction,” *Phys. Rev. D* **83**, 032008 (2011) [arXiv:1004.4855 [hep-ex]].
216. T. Aaltonen *et al.* [CDF Collaboration], “Search for R-parity Violating Decays of  $\tau$  sneutrinos to  $e\mu$ ,  $\mu\tau$ , and  $e\tau$  Pairs in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96$  TeV,” *Phys. Rev. Lett.* **105**, 191801 (2010) [arXiv:1004.3042 [hep-ex]].
217. T. Aaltonen *et al.* [CDF Collaboration], “First Measurement of the Ratio  $\sigma(t\bar{t})/\sigma(Z/\gamma^* \rightarrow \ell\bar{\ell})$  and Precise Extraction of the  $t\bar{t}$  Cross Section,” *Phys. Rev. Lett.* **105**, 012001 (2010) [arXiv:1004.3224 [hep-ex]].
218. M. Anelli, S. Bertolucci, C. Bini, P. Branchini, G. Corradi, C. Curceanu, G. De Zorzi and A. Di Domenico *et al.*, “Measurement of the neutron detection efficiency of a 80arXiv:1004.2241 [physics.ins-det].
219. T. Aaltonen *et al.* [CDF Collaboration], “Observation of Single Top Quark Production and Measurement of  $-\text{V}t\text{b}$ — with CDF,” *Phys. Rev. D* **82**, 112005 (2010) [arXiv:1004.1181 [hep-ex]].
220. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of Z gamma Production in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96$  TeV,” *Phys. Rev. D* **82**, 031103 (2010) [arXiv:1004.1140 [hep-ex]].
221. G. Amelino-Camelia, F. Archilli, D. Babusci, D. Badoni, G. Bencivenni, J. Bernabeu, R. A. Bertlmann and D. R. Boito *et al.*, “Physics with the KLOE-2 experiment at the upgraded DAΦNE,” *Eur. Phys. J. C* **68**, 619 (2010) [arXiv:1003.3868 [hep-ex]].
222. T. Aaltonen *et al.* [CDF Collaboration], “Studying the Underlying Event in Drell-Yan and High Transverse Momentum Jet Production at the Tevatron,” *Phys. Rev. D* **82**, 034001 (2010) [arXiv:1003.3146 [hep-ex]].
223. T. A. Aaltonen *et al.* [CDF Collaboration], “Measurement of  $d\sigma/dy$  of Drell-Yan  $e^+e^-$  pairs in the  $Z$  Mass Region from  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96$  TeV,” *Phys. Lett. B* **692**, 232 (2010) [arXiv:0908.3914 [hep-ex], arXiv:0908.3914 [hep-ex]].
224. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of W-Boson Polarization in Top-quark Decay in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96$  TeV,” *Phys. Rev. Lett.* **105**, 042002 (2010) [arXiv:1003.0224 [hep-ex]].

225. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the  $t\bar{t}$  Production Cross Section in  $p\bar{p}$  Collisions at  $\sqrt{s}=1.96$  TeV using Soft Electron b-Tagging,” *Phys. Rev. D* **81**, 092002 (2010) [arXiv:1002.3783 [hep-ex]].
226. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the Top Pair Production Cross Section in the Dilepton Decay Channel in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96$  TeV,” *Phys. Rev. D* **82**, 052002 (2010) [arXiv:1002.2919 [hep-ex]].
227. K. : F. Archilli *et al.* [2 Collaboration], “Technical Design Report of the Inner Tracker for the KLOE-2 experiment,” arXiv:1002.2572 [physics.ins-det].
228. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the Top Quark Mass and  $p\bar{p} \rightarrow t\bar{t}$  Cross Section in the All-Hadronic Mode with the CDFII Detector,” *Phys. Rev. D* **81**, 052011 (2010) [arXiv:1002.0365 [hep-ex]].
229. T. Aaltonen *et al.* [CDF Collaboration], “Search for single top quark production in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV in the missing transverse energy plus jets topology,” *Phys. Rev. D* **81**, 072003 (2010) [arXiv:1001.4577 [hep-ex]].
230. T. Aaltonen *et al.* [CDF and D0 Collaborations], “Combination of Tevatron searches for the standard model Higgs boson in the  $W+W-$  decay mode,” *Phys. Rev. Lett.* **104**, 061802 (2010) [arXiv:1001.4162 [hep-ex]].
231. T. Aaltonen *et al.* [CDF Collaboration], “Inclusive Search for Standard Model Higgs Boson Production in the  $WW$  Decay Channel using the CDF II Detector,” *Phys. Rev. Lett.* **104**, 061803 (2010) [arXiv:1001.4468 [hep-ex]].
232. S. Bethke *et al.* [CDF Collaboration], “Comments and reply on: ‘Study of multi-muon events produced in interactions at  $\sqrt{s} = 1.96$  TeV’,” *Eur. Phys. J. C* **68**, 119 (2010).
233. T. Aaltonen *et al.* [CDF Collaboration], “Search for New Physics with a Dijet plus Missing Transverse Energy Signature in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96$  TeV,” *Phys. Rev. Lett.* **105**, 131801 (2010) [arXiv:0912.4691 [hep-ex]].
234. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the  $W+ W-$  Production Cross Section and Search for Anomalous  $WW$  gamma and  $WWZ$  Couplings in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96$ -TeV,” *Phys. Rev. Lett.* **104**, 201801 (2010) [arXiv:0912.4500 [hep-ex]].
235. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the  $\Lambda(1520)$  Lifetime in  $\Lambda(1520) \rightarrow \Lambda^0 \pi^0$  Decays in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96$ -TeV,” *Phys. Rev. Lett.* **104**, 102002 (2010) [arXiv:0912.3566 [hep-ex]].
236. T. Aaltonen *et al.* [CDF Collaboration], “A Study of the associated production of photons and b-quark jets in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$ -TeV,” *Phys. Rev. D* **81**, 052006 (2010) [arXiv:0912.3453 [hep-ex]].

237. T. Aaltonen *et al.* [CDF Collaboration], “Search for Technicolor Particles Produced in Association with a W Boson at CDF,” *Phys. Rev. Lett.* **104**, 111802 (2010) [arXiv:0912.2059 [hep-ex]].
238. T. Aaltonen *et al.* [CDF Collaboration], “Search for Pair Production of Supersymmetric Top Quarks in Dilepton Events from p anti-p Collisions at  $S^{*(1/2)} = 1.96$  TeV,” *Phys. Rev. Lett.* **104**, 251801 (2010) [arXiv:0912.1308 [hep-ex]].
239. T. Aaltonen *et al.* [CDF Collaboration], “Search for New Bottomlike Quark Pair Decays  $Q Q\text{-bar} \rightarrow j$  ( $t W\text{-}$ ) ( $t\text{-bar} W\text{+}$ ) in Same-Charge Dilepton Events,” *Phys. Rev. Lett.* **104**, 091801 (2010) [arXiv:0912.1057 [hep-ex]].
240. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the WW+WZ Production Cross Section Using the Lepton+Jets Final State at CDF II,” *Phys. Rev. Lett.* **104**, 101801 (2010) [arXiv:0911.4449 [hep-ex]].
241. T. Aaltonen *et al.* [CDF Collaboration], “A Search for the Higgs Boson Using Neural Networks in Events with Missing Energy and b-quark Jets in p anti-p Collisions at  $s^{*(1/2)} = 1.96$ -TeV,” *Phys. Rev. Lett.* **104**, 141801 (2010) [arXiv:0911.3935 [hep-ex]].
242. T. Aaltonen *et al.* [CDF Collaboration], “Search for New Color-Octet Vector Particle Decaying to t anti-t in p anti-p Collisions at  $s^{*(1/2)} = 1.96$ -TeV,” *Phys. Lett. B* **691**, 183 (2010) [arXiv:0911.3112 [hep-ex]].
243. T. Aaltonen *et al.* [CDF Collaboration], “Top Quark Mass Measurement using mT2 in the Dilepton Channel at CDF,” *Phys. Rev. D* **81**, 031102 (2010) [arXiv:0911.2956 [hep-ex]].
244. T. Aaltonen *et al.* [CDF Collaboration], “Measurements of branching fraction ratios and CP asymmetries in  $B^{+-} \rightarrow j D(\text{CP}) K^{+-}$  decays in hadron collisions,” *Phys. Rev. D* **81**, 031105 (2010) [arXiv:0911.0425 [hep-ex]].
245. T. Aaltonen *et al.* [CDF Collaboration], “Search for Anomalous Production of Events with Two Photons and Additional Energetic Objects at CDF,” *Phys. Rev. D* **82**, 052005 (2010) [arXiv:0910.5170 [hep-ex]].
246. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the Inclusive Isolated Prompt Photon Cross Section in p anti-p Collisions at  $s^{*(1/2)} = 1.96$ -TeV using the CDF Detector,” *Phys. Rev. D* **80**, 111106 (2009) [arXiv:0910.3623 [hep-ex]].
247. T. Aaltonen *et al.* [CDF Collaboration], “Search for Supersymmetry with Gauge-Mediated Breaking in Diphoton Events with Missing Transverse Energy at CDF II,” *Phys. Rev. Lett.* **104**, 011801 (2010) [arXiv:0910.3606 [hep-ex]].
248. T. Aaltonen *et al.* [CDF Collaboration], “Measurements of the top-quark mass using charged particle tracking,” *Phys. Rev. D* **81**, 032002 (2010) [arXiv:0910.0969 [hep-ex]].

249. M. Cordelli, G. Corradi, F. Happacher, M. Martini, S. Miscetti, C. Paglia, A. Saputi and I. Sarra *et al.*, “QCALT: A tile calorimeter for KLOE-2 experiment,” In \*Como 2009, Astroparticle, particle and space physics, detectors and medical physics applications\* 404-408
250. M. Cordelli, F. Happacher, M. Martini, S. Miscetti, I. Sarra, M. Schioppa, S. Stucci and G. Xu, “Test of a LYSO matrix with an electron beam between 100-MeV and 500-MeV for KLOE-2,” In \*Como 2009, Astroparticle, particle and space physics, detectors and medical physics applications\* 372-376
251. T. Aaltonen *et al.* [CDF Collaboration], “First Measurement of the b-jet Cross Section in Events with a W Boson in p anti-p Collisions at  $s^{*}(1/2) = 1.96$ -TeV,” Phys. Rev. Lett. **104**, 131801 (2010) [arXiv:0909.1505 [hep-ex]].
252. T. Aaltonen *et al.* [CDF Collaboration], “A Search for the Higgs Boson Produced in Association with  $Z \rightarrow \ell^+\ell^-$  Using the Matrix Element Method at CDF II,” Phys. Rev. D **80**, 071101 (2009) [arXiv:0908.3534 [hep-ex]].
253. T. Aaltonen *et al.* [CDF Collaboration], “Search for charged Higgs bosons in decays of top quarks in p anti-p collisions at  $s^{*}(1/2) = 1.96$  TeV,” Phys. Rev. Lett. **103**, 101803 (2009) [arXiv:0907.1269 [hep-ex]].
254. T. Aaltonen *et al.* [CDF Collaboration], “A Search for the Associated Production of the Standard-Model Higgs Boson in the All-Hadronic Channel,” Phys. Rev. Lett. **103**, 221801 (2009) [arXiv:0907.0810 [hep-ex]].
255. T. Aaltonen *et al.* [CDF Collaboration], “Search for a Higgs Boson in  $WH \rightarrow \ell\nu b\bar{b}$  in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96$  TeV,” Phys. Rev. Lett. **103**, 101802 (2009) [arXiv:0906.5613 [hep-ex]].
256. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the Top Quark Mass Using the Invariant Mass of Lepton Pairs in Soft Muon b-tagged Events,” Phys. Rev. D **80**, 051104 (2009) [arXiv:0906.5371 [hep-ex]].
257. T. Aaltonen *et al.* [CDF Collaboration], “Precision Measurement of the X(3872) Mass in J/psi pi+ pi- Decays,” Phys. Rev. Lett. **103**, 152001 (2009) [arXiv:0906.5218 [hep-ex]].
258. M. Anelli, G. Battistoni, S. Bertolucci, C. Bini, P. Branchini, C. Curceanu, G. De Zorzi and A. Domenico *et al.*, “Measurement and simulation of the neutron detection efficiency with a Pb-scintillating fiber calorimeter,” J. Phys. Conf. Ser. **160**, 012023 (2009).
259. F. Happacher, M. Martini, S. Miscetti and I. Sarra, “Tile and crystal calorimeter for KLOE-2 experiment,” Nucl. Phys. Proc. Suppl. **197**, 215 (2009) [arXiv:0906.1157 [physics.ins-det]].

260. M. Cordelli, F. Happacher, M. Martini, S. Miscetti, I. Sarra, M. Schioppa, S. Stucci and G. Xu, “Test of a LYSO matrix with an electron beam between 100-MeV and 500-MeV for KLOE-2,” Nucl. Instrum. Meth. A **617**, 109 (2010) [arXiv:0906.1144 [physics.ins-det]].
261. M. Anelli, S. Bertolucci, C. Bini, P. Branchini, G. Corradi, C. Curceanu, G. De Zorzi and A. Di Domenico *et al.*, “Measurement of neutron detection efficiency between 22-MeV and 174- MeV using two different kinds of Pb-scintillating fiber sampling calorimeters,” Nucl. Instrum. Meth. A **617**, 107 (2010) [arXiv:0906.1143 [physics.ins-det]].
262. M. Cordelli, G. Corradi, F. Happacher, M. Martini, S. Miscetti, C. Paglia, A. Saputi and I. Sarra *et al.*, “QCALT: A Tile calorimeter for KLOE-2 experiment,” Nucl. Instrum. Meth. A **617**, 105 (2010) [arXiv:0906.1133 [physics.ins-det]].
263. T. Aaltonen *et al.* [CDF Collaboration], “Search for Higgs bosons predicted in two-Higgs-doublet models via decays to tau lepton pairs in 1.96-TeV p anti-p collisions,” Phys. Rev. Lett. **103**, 201801 (2009) [arXiv:0906.1014 [hep-ex]].
264. T. Aaltonen *et al.* [CDF Collaboration], “Searching the Inclusive Lepton + Photon + Missing E(T) + b-quark Signature for Radiative Top Quark Decay and Non-Standard-Model Processes,” Phys. Rev. D **80**, 011102 (2009) [arXiv:0906.0518 [hep-ex]].
265. T. Aaltonen *et al.* [CDF Collaboration], “First Observation of Vector Boson Pairs in a Hadronic Final State at the Tevatron Collider,” Phys. Rev. Lett. **103**, 091803 (2009) [arXiv:0905.4714 [hep-ex]].
266. T. Aaltonen *et al.* [CDF Collaboration], “Search for Standard Model Higgs Boson Production in Association with a W Boson using a Neural Network Discriminant at CDF,” Phys. Rev. D **80**, 012002 (2009) [arXiv:0905.3155 [hep-ex]].
267. T. Aaltonen *et al.* [CDF Collaboration], “Observation of the Omega(b)- Baryon and Measurement of the Properties of the Xi(b)- and Omega(b)- Baryons,” Phys. Rev. D **80**, 072003 (2009) [arXiv:0905.3123 [hep-ex]].
268. T. Aaltonen *et al.* [CDF Collaboration], “Production of psi(2S) Mesons in p anti-p Collisions at 1.96-TeV,” Phys. Rev. D **80**, 031103 (2009) [arXiv:0905.1982 [hep-ex]].
269. T. Aaltonen *et al.* [CDF Collaboration], “Search for a Fermiophobic Higgs Boson Decaying into Diphotons in p anti-p Collisions at  $s^{*}(1/2) = 1.96\text{-TeV}$ ,” Phys. Rev. Lett. **103**, 061803 (2009) [arXiv:0905.0413 [hep-ex]].
270. T. Aaltonen *et al.* [CDF Collaboration], “Search for the Neutral Current Top Quark Decay  $t \rightarrow j Z c$  Using Ratio of Z-Boson + 4 Jets to W-Boson + 4 Jets Production,” Phys. Rev. D **80**, 052001 (2009) [arXiv:0905.0277 [hep-ex]].

271. T. Aaltonen *et al.* [CDF Collaboration], “Search for Anomalous Production of Events with a Photon, Jet, b-quark Jet, and Missing Transverse Energy,” *Phys. Rev. D* **80**, 052003 (2009) [arXiv:0905.0231 [hep-ex]].
272. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of Particle Production and Inclusive Differential Cross Sections in p anti-p Collisions at  $s^{**}(1/2) = 1.96$ -TeV,” *Phys. Rev. D* **79**, 112005 (2009) [Erratum-ibid. *D* **82**, 119903 (2010)] [arXiv:0904.1098 [hep-ex]].
273. T. Aaltonen *et al.* [CDF Collaboration], “A Measurement of the t anti-t Cross Section in p anti-p Collisions at  $s^{**}(1/2) = 1.96$ -TeV using Dilepton Events with a Lepton plus Track Selection,” *Phys. Rev. D* **79**, 112007 (2009) [arXiv:0903.5263 [hep-ex]].
274. T. Aaltonen *et al.* [CDF Collaboration], “First Measurement of the t anti-t Differential Cross Section  $d\sigma/dM(t\text{ anti-t})$  in p anti-p Collisions at  $s^{**}(1/2)=1.96$ -TeV,” *Phys. Rev. Lett.* **102**, 222003 (2009) [arXiv:0903.2850 [hep-ex]].
275. T. Aaltonen *et al.* [CDF Collaboration], “Search for Gluino-Mediated Sbottom Production in p anti-p Collisions at  $s^{**}(1/2) = 1.96$ -TeV,” *Phys. Rev. Lett.* **102**, 221801 (2009) [arXiv:0903.2618 [hep-ex]].
276. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the b-Hadron Production Cross Section Using Decays to MU- d0 X Final States in p anti-p Collisions at  $s^{**}(1/2) = 1.96$ -TeV,” *Phys. Rev. D* **79**, 092003 (2009) [arXiv:0903.2403 [hep-ex]].
277. T. Aaltonen *et al.* [CDF Collaboration], “Evidence for a Narrow Near-Threshold Structure in the  $J/\psi\phi$  Mass Spectrum in  $B^+ \rightarrow J/\psi\phi K^+$  Decays,” *Phys. Rev. Lett.* **102**, 242002 (2009) [arXiv:0903.2229 [hep-ex]].
278. T. Aaltonen *et al.* [CDF Collaboration], “Search for narrow resonances lighter than  $\Upsilon$  mesons,” *Eur. Phys. J. C* **62**, 319 (2009) [arXiv:0903.2060 [hep-ex]].
279. T. Aaltonen *et al.* [CDF Collaboration], “First Observation of Electroweak Single Top Quark Production,” *Phys. Rev. Lett.* **103**, 092002 (2009) [arXiv:0903.0885 [hep-ex]].
280. T. Aaltonen *et al.* [CDF Collaboration], “Search for WW and WZ production in lepton plus jets final state at CDF,” *Phys. Rev. D* **79**, 112011 (2009) [arXiv:0903.0814 [hep-ex]].
281. T. Aaltonen *et al.* [CDF Collaboration], “Search for the Production of Narrow t anti-b Resonances in 1.9 fb<sup>-1</sup> of p anti-p Collisions at  $s^{**}(1/2) = 1.96$ -TeV,” *Phys. Rev. Lett.* **103**, 041801 (2009) [arXiv:0902.3276 [hep-ex]].

282. T. Aaltonen *et al.* [CDF Collaboration], “Search for exclusive  $Z$  boson production and observation of high mass  $p\bar{p} \rightarrow \gamma\gamma \rightarrow p + \ell\ell + \bar{p}$  events in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV,” Phys. Rev. Lett. **102**, 222002 (2009) [arXiv:0902.2816 [hep-ex]].
283. T. Aaltonen *et al.* [CDF Collaboration], “Observation of exclusive charmonium production and  $\gamma\gamma \rightarrow \mu^+\mu^-$  in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.96$ -TeV,” Phys. Rev. Lett. **102**, 242001 (2009) [arXiv:0902.1271 [hep-ex]].
284. T. Aaltonen *et al.* [CDF Collaboration], “Search for Long-Lived Massive Charged Particles in 1.96 TeV  $p\bar{p}$  Collisions,” Phys. Rev. Lett. **103**, 021802 (2009) [arXiv:0902.1266 [hep-ex]].
285. M. Anelli *et al.* [KLOE Collaboration], “Measurement of the detection efficiency of the KLOE calorimeter for neutrons between 22-MeV and 174-MeV,” Nucl. Instrum. Meth. A **598**, 244 (2009).
286. R. M. Carey, K. R. Lynch, J. P. Miller, B. L. Roberts, W. M. Morse, Y. K. Semertzides, V. P. Druzhinin and B. I. Khazin *et al.*, “The New (g-2) Experiment: A proposal to measure the muon anomalous magnetic moment to  $\pm 0.14$  ppm precision,” FERMILAB-PROPOSAL-0989.
287. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the  $t\bar{t}$  Production Cross Section in  $2 \text{ fb}^{-1}$  of  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96$  TeV Using Lepton Plus Jets Events with Soft Muon b-Tagging,” Phys. Rev. D **79**, 052007 (2009) [arXiv:0901.4142 [hep-ex]].
288. T. Aaltonen *et al.* [CDF Collaboration], “Search for the Decays  $B_s^0 \rightarrow e^+\mu^-$  and  $B_s^0 \rightarrow e^+e^-$  in CDF Run II,” Phys. Rev. Lett. **102**, 201801 (2009) [arXiv:0901.3803 [hep-ex]].
289. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the top quark mass at CDF using the ‘neutrino  $\phi$  weighting’ template method on a lepton plus isolated track sample,” Phys. Rev. D **79**, 072005 (2009) [arXiv:0901.3773 [hep-ex]].
290. T. Aaltonen *et al.* [CDF Collaboration], “Direct Measurement of the  $W$  Production Charge Asymmetry in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96$  TeV,” Phys. Rev. Lett. **102**, 181801 (2009) [arXiv:0901.2169 [hep-ex]].
291. F. Happacher [CDF Collaboration], “Multi-muon events at CDF,” PoS EPS **-HEP2009**, 247 (2009).
292. P. Branchini *et al.* [KLOE2 Collaboration], “Detector upgrade for the KLOE2 experiment: The calorimeter system,” PoS KAON **09**, 013 (2009).
293. F. Happacher [CDF Collaboration], “Study of multi-muon events at CDF,” Nuovo Cim. C **32N5-6**, 371 (2009).

294. T. Aaltonen *et al.* [CDF Collaboration], “Top Quark Mass Measurement in the Lepton plus Jets Channel Using a Modified Matrix Element Method,” Phys. Rev. D **79**, 072001 (2009) [arXiv:0812.4469 [hep-ex]].
295. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of Cross Sections for  $b$  Jet Production in Events with a  $Z$  Boson in  $p^-$  anti- $p$  Collisions at  $\sqrt{s} = 1.96$ -TeV,” Phys. Rev. D **79**, 052008 (2009) [arXiv:0812.4458 [hep-ex]].
296. T. Aaltonen *et al.* [CDF Collaboration], “Observation of New Charmless Decays of Bottom Hadrons,” Phys. Rev. Lett. **103**, 031801 (2009) [arXiv:0812.4271 [hep-ex]].
297. T. Aaltonen *et al.* [CDF Collaboration], “Search for new particles decaying into dijets in proton-antiproton collisions at  $s^{*(1/2)} = 1.96$ -TeV,” Phys. Rev. D **79**, 112002 (2009) [arXiv:0812.4036 [hep-ex]].
298. T. Aaltonen *et al.* [CDF Collaboration], “Search for top-quark production via flavor-changing neutral currents in  $W+1$  jet events at CDF,” Phys. Rev. Lett. **102**, 151801 (2009) [arXiv:0812.3400 [hep-ex]].
299. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the  $k_T$  Distribution of Particles in Jets Produced in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96$ -TeV,” Phys. Rev. Lett. **102**, 232002 (2009) [arXiv:0811.2820 [hep-ex]].
300. T. Aaltonen *et al.* [CDF Collaboration], “Inclusive Search for Squark and Gluino Production in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96$ -TeV,” Phys. Rev. Lett. **102**, 121801 (2009) [arXiv:0811.2512 [hep-ex]].
301. T. Aaltonen *et al.* [CDF Collaboration], “Top Quark Mass Measurement in the  $t\bar{t}$  All Hadronic Channel using a Matrix Element Technique in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96$ -TeV,” Phys. Rev. D **79**, 072010 (2009) [arXiv:0811.1062 [hep-ex]].
302. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of W-Boson Helicity Fractions in Top-Quark Decays Using  $\cos\theta^*$ ,” Phys. Lett. B **674**, 160 (2009) [arXiv:0811.0344 [hep-ex]].
303. T. Aaltonen *et al.* [CDF Collaboration], “A Search for high-mass resonances decaying to dimuons at CDF,” Phys. Rev. Lett. **102**, 091805 (2009) [arXiv:0811.0053 [hep-ex]].
304. T. Aaltonen *et al.* [CDF Collaboration], “Study of multi-muon events produced in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$ -TeV,” arXiv:0810.5357 [hep-ex].
305. T. Aaltonen *et al.* [CDF Collaboration], “Search for new physics in the  $\mu\mu + e/\mu + \cancel{E}_T$  channel with a low- $p_T$  lepton threshold at the Collider Detector at Fermilab,” Phys. Rev. D **79**, 052004 (2009) [arXiv:0810.3522 [hep-ex]].
306. T. Aaltonen *et al.* [CDF Collaboration], Phys. Rev. D **79**, 032001 (2009) [arXiv:0810.3213 [hep-ex]].

307. T. Aaltonen *et al.* [CDF Collaboration], Phys. Rev. Lett. **102**, 031801 (2009) [arXiv:0810.2059 [hep-ex]].
308. P. Giromini, F. Happacher, M. J. Kim, M. Kruse, K. Pitts, F. Ptohos and S. Torre, [arXiv:0810.5730 [hep-ph]].
309. T. Aaltonen *et al.* [CDF Collaboration], Phys. Rev. Lett. **102**, 102003 (2009) [arXiv:0809.5007 [hep-ex]].
310. T. Aaltonen *et al.* [CDF Collaboration], “Search for Maximal Flavor Violating Scalars in Same-Charge Lepton Pairs in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96$  TeV,” arXiv:0809.4903 [hep-ex].
311. T. Aaltonen *et al.* [CDF Collaboration], “The first measurement of the top quark mass at CDF II in the lepton+jets and dilepton channels simultaneously,” arXiv:0809.4808 [hep-ex].
312. T. Aaltonen *et al.* [CDF Collaboration], “Search for a Higgs Boson Decaying to Two W Bosons at CDF,” arXiv:0809.3930 [hep-ex].
313. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the Single Top Quark Production Cross Section at CDF,” arXiv:0809.2581 [hep-ex].
314. T. Aaltonen *et al.* [CDF Collaboration], “Search for Supersymmetry in  $p\bar{p}$  Collisions at  $\sqrt{s}=1.96$  TeV Using the Trilepton Signature of Chargino-Neutralino Production,” arXiv:0808.2446 [hep-ex].
315. T. Aaltonen *et al.* [CDF Collaboration], “First Direct Bound on the Total Width of the Top Quark in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96$  TeV,” arXiv:0808.2167 [hep-ex].
316. T. Aaltonen *et al.* [The CDF Collaboration], “Search for Doubly Charged Higgs Bosons with Lepton-Flavor-Violating Decays involving Tau Leptons,” arXiv:0808.2161 [hep-ex].
317. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the top quark mass with dilepton events selected using neuroevolution at CDF,” arXiv:0807.4652 [hep-ex].
318. T. Aaltonen *et al.* [CDF Collaboration], “Search for the Higgs boson produced with  $Z \rightarrow \ell^+\ell^-$  in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV,” arXiv:0807.4493 [hep-ex].
319. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the fraction of t-tbar production via gluon-gluon fusion in p-pbar collisions at  $\sqrt{s}=1.96$  TeV,” arXiv:0807.4262 [hep-ex].
320. T. Aaltonen *et al.* [CDF Collaboration], “Search for large extra dimensions in final states containing one photon or jet and large missing transverse energy produced in p-pbar collisions at  $\sqrt{s} = 1.96$  TeV,” arXiv:0807.3132 [hep-ex].

321. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the Inclusive Jet Cross Section at the Fermilab Tevatron p-pbar Collider Using a Cone-Based Jet Algorithm,” arXiv:0807.2204 [hep-ex].
322. T. Aaltonen *et al.* [CDF Collaboration], “Forward-Backward Asymmetry in Top Quark Production in ppbar Collisions at  $\sqrt{s}=1.96$  TeV,” arXiv:0806.2472 [hep-ex].
323. T. Aaltonen *et al.* [CDF Collaboration], “Search for the Rare Decays  $B^{+-} \rightarrow \mu^+\mu^-K^+, B^0 \rightarrow \mu^+\mu^-K^0(892)$ , and  $B_s^0 \rightarrow \mu^+\mu^- \phi$  at CDF,” arXiv:0804.3908 [hep-ex].
324. T. Aaltonen *et al.* [CDF Collaboration], “Search for Heavy, Long-Lived Neutralinos that Decay to Photons at CDF II Using Photon Timing,” Phys. Rev. D **78** (2008) 032015 [arXiv:0804.1043 [hep-ex]].
325. T. Aaltonen *et al.* [CDF Collaboration], “Search for Hadronic Decays of W and Z Bosons in Photon Events in ppbar Collisions at  $\sqrt{s}=1.96$  GeV,” arXiv:0803.4264 [hep-ex].
326. T. Aaltonen *et al.* [CDF Collaboration], “Search for Standard Model Higgs Boson Production in Association with a W Boson at CDF,” Phys. Rev. D **78** (2008) 032008 [arXiv:0803.3493 [hep-ex]].
327. T. Aaltonen *et al.* [CDF Collaboration], “Search for Pair Production of Scalar Top Quarks Decaying to a tau Lepton and a b Quark in ppbar Collisions at  $\sqrt{s}=1.96$  TeV,” Phys. Rev. Lett. **101** (2008) 071802 [arXiv:0802.3887 [hep-ex]].
328. M. Anelli *et al.*, “Measurement and simulation of the neutron response and detection efficiency of a Pb-scintillating fiber calorimeter,” Nucl. Instrum. Meth. A **581** (2007) 368 [IEEE Trans. Nucl. Sci. **55** (2008) 1409].
329. T. Aaltonen *et al.* [CDF Collaboration], “Two-Particle Momentum Correlations in Jets Produced in ppbar Collisions at  $\sqrt{s}=1.96$ -TeV,” arXiv:0802.3182 [hep-ex].
330. T. Aaltonen *et al.* [CDF Collaboration], “Search for the Higgs boson in events with missing transverse energy and b quark jets produced in proton-antiproton collisions at  $\sqrt{s}=1.96$  TeV,” Phys. Rev. Lett. **100** (2008) 211801 [arXiv:0802.0432 [hep-ex]].
331. T. Aaltonen *et al.* [CDF Collaboration], “First Measurement of ZZ Production in ppbar Collisions at  $\sqrt{s}=1.96$  TeV,” Phys. Rev. Lett. **100** (2008) 201801 [arXiv:0801.4806 [hep-ex]].
332. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of Ratios of Fragmentation Fractions for Bottom Hadrons in p-pbar Collisions at  $\sqrt{s}=1.96$  TeV,” Phys. Rev. D **77** (2008) 072003 [arXiv:0801.4375 [hep-ex]].

333. T. Aaltonen *et al.* [CDF Collaboration], “Search for Heavy Top-like Quarks  $t'$   $\rightarrow$   $Wq$  Using Lepton Plus Jets Events in 1.96 TeV Proton-Antiproton Collisions,” Phys. Rev. Lett. **100** (2008) 161803 [arXiv:0801.3877 [hep-ex]].
334. T. Aaltonen *et al.* [CDF Collaboration], “Search for New Heavy Particles Decaying to  $Z^0 Z^0 \rightarrow eeee$  in p-pbar Collisions at  $\sqrt{s} = 1.96$  TeV,” Phys. Rev. D **78** (2008) 012008 [arXiv:0801.1129 [hep-ex]].
335. T. Aaltonen *et al.* [CDF Collaboration], “First Measurement of the Fraction of Top Quark Pair Production Through Gluon-Gluon Fusion,” arXiv:0712.3273 [hep-ex].
336. T. Aaltonen *et al.* [CDF Collaboration], “Model-Independent Global Search for New High-pT Physics at CDF,” arXiv:0712.2534 [hep-ex].
337. T. Aaltonen *et al.* [CDF Collaboration], “First Flavor-Tagged Determination of Bounds on Mixing-Induced CP Violation in  $B_s \rightarrow J/\psi \phi$  Decays,” Phys. Rev. Lett. **100** (2008) 161802 [arXiv:0712.2397 [hep-ex]].
338. T. Aaltonen *et al.* [CDF collaboration], “Measurement of Lifetime and Decay-Width Difference in  $B_0 \rightarrow J/\psi \phi$  Decays,” Phys. Rev. Lett. **100** (2008) 121803 [arXiv:0712.2348 [hep-ex]].
339. T. Aaltonen *et al.* [CDF Collaboration], “Evidence for  $D_0$ - $D_0$ bar mixing using the CDF II Detector,” Phys. Rev. Lett. **100** (2008) 121802 [arXiv:0712.1567 [hep-ex]].
340. T. Aaltonen *et al.* [CDF Collaboration], “Search for  $B_s \rightarrow \mu^+\mu^-$  and  $B_d \rightarrow \mu^+\mu^-$  Decays with 2fb-1 of ppbar Collisions,” Phys. Rev. Lett. **100** (2008) 101802 [arXiv:0712.1708 [hep-ex]].
341. T. Aaltonen *et al.* [CDF Collaboration], “Observation of the Decay  $B_c^\pm \rightarrow J/\psi \pi^\pm$  and Measurement of the  $B_c^\pm$  Mass,” Phys. Rev. Lett. **100** (2008) 182002 [arXiv:0712.1506 [hep-ex]].
342. T. Aaltonen *et al.* [CDF Collaboration], “Model-Independent and Quasi-Model-Independent Search for New Physics at CDF,” arXiv:0712.1311 [hep-ex].
343. T. Aaltonen *et al.* [CDF Run II Collaboration], “Observation of Exclusive Dijet Production at the Fermilab Tevatron p-pbar Collider,” Phys. Rev. D **77** (2008) 052004 [arXiv:0712.0604 [hep-ex]].
344. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the cross section for W-boson production in association with jets in ppbar collisions at  $\sqrt{s} = 1.96$  TeV,” Phys. Rev. D **77** (2008) 011108 [arXiv:0711.4044 [hep-ex]].
345. T. Aaltonen *et al.* [CDF - Run II Collaboration], “Measurement of Inclusive Jet Cross Sections in  $Z/g^{**}(\rightarrow ee) +$  jets Production in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96$  TeV,” Phys. Rev. Lett. **100** (2008) 102001 [arXiv:0711.3717 [hep-ex]].

346. T. Aaltonen *et al.* [CDF Collaboration], “Search for chargino-neutralino production in p-pbar collisions at 1.96 TeV with high pT leptons,” Phys. Rev. D **77** (2008) 052002 [arXiv:0711.3161 [hep-ex]].
347. T. Aaltonen *et al.* [CDF Collaboration], “First Measurement of the Production of a W Boson in Association with a Single Charm Quark in Proton Anti-proton Collisions at sqrt(s)=1.96 TeV,” Phys. Rev. Lett. **100** (2008) 091803 [arXiv:0711.2901 [hep-ex]].
348. T. Aaltonen *et al.* [CDF Collaboration], “Limits on the Production of Narrow t-tbar Resonances in p-pbar Collisions at sqrt(s)=1.96 TeV,” Phys. Rev. D **77** (2008) 051102 [arXiv:0710.5335 [hep-ex]].
349. T. Aaltonen *et al.* [CDF Collaboration], “Search for Standard Model Higgs Bosons Produced in Association with W Bosons,” Phys. Rev. Lett. **100** (2008) 041801 [arXiv:0710.4363 [hep-ex]].
350. T. Aaltonen *et al.* [CDF Collaboration], “Observation of Orbitally Excited  $B_s$  Mesons,” Phys. Rev. Lett. **100** (2008) 082001 [arXiv:0710.4199 [hep-ex]].
351. T. Aaltonen *et al.* [CDF Collaboration], “A direct Measurement of the W Boson Width in ppbar Collisions at sqrt(s) = 1.96 TeV,” Phys. Rev. Lett. **100** (2008) 071801 [arXiv:0710.4112 [hep-ex]].
352. T. Aaltonen *et al.* [CDF Collaboration], “Cross Section Constrained Top Quark Mass Measurement from Dilepton Events at the Tevatron,” Phys. Rev. Lett. **100** (2008) 062005 [arXiv:0710.4037 [hep-ex]].
353. T. Aaltonen *et al.* [CDF Run II Collaboration], “First observation of the decay  $B_s^0 \rightarrow D_s^- D_s^+$  and measurement of its branching ratio,” Phys. Rev. Lett. **100** (2008) 021803.
354. T. Aaltonen *et al.* [CDF Run II Collaboration], “Measurement of correlated b-bbar production in p-pbar collisions at s\*\*(1/2) = 1960 GeV,” Phys. Rev. D **77** (2008) 072004 [arXiv:0710.1895 [hep-ex]].
355. T. Aaltonen *et al.* [CDF Collaboration], “Search for resonant  $t\bar{t}$  production in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$ -TeV,” Phys. Rev. Lett. **100** (2008) 231801 [arXiv:0709.0705 [hep-ex]].
356. T. Aaltonen *et al.* [CDF Collaboration], “First Run II Measurement of the W Boson Mass,” Phys. Rev. D **77** (2008) 112001 [arXiv:0708.3642 [hep-ex]].
357. T. Aaltonen *et al.* [CDF Collaboration], “Search for Direct Pair Production of Supersymmetric Top and Supersymmetric Bottom Quarks in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96$ -TeV,” Phys. Rev. D **76** (2007) 072010 [arXiv:0707.2567 [hep-ex]].

358. T. Aaltonen *et al.* [CDF Collaboration], “Search for new physics in high mass electron-positron events in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96\text{-TeV}$ ,” Phys. Rev. Lett. **99** (2007) 171802 [arXiv:0707.2524 [hep-ex]].
359. T. Aaltonen *et al.* [CDF Collaboration], “Search for exclusive  $\gamma\gamma$  production in hadron-hadron collisions,” Phys. Rev. Lett. **99** (2007) 242002 [arXiv:0707.2374 [hep-ex]].
360. T. Aaltonen *et al.* [CDF Collaboration], “Search for chargino-neutralino production in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96\text{-TeV}$ ,” Phys. Rev. Lett. **99** (2007) 191806 [arXiv:0707.2362 [hep-ex]].
361. T. Aaltonen *et al.* [CDF Collaboration], “Search for a high-mass diphoton state and limits on Randall-Sundrum gravitons at CDF,” Phys. Rev. Lett. **99** (2007) 171801 [arXiv:0707.2294 [hep-ex]].
362. T. Aaltonen *et al.* [CDF Collaboration], “Observation and mass measurement of the baryon  $Xi_b^-$ ,” Phys. Rev. Lett. **99** (2007) 052002 [arXiv:0707.0589 [hep-ex]].
363. T. Aaltonen *et al.* [CDF Collaboration], “First Measurement of the W Boson Mass in Run II of the Tevatron,” Phys. Rev. Lett. **99** (2007) 151801 [arXiv:0707.0085 [hep-ex]].
364. T. Aaltonen *et al.* [CDF Collaboration], “First observation of heavy baryons  $\Sigma_b$  and  $\Sigma_b^*$ ,” Phys. Rev. Lett. **99** (2007) 202001 [arXiv:0706.3868 [hep-ex]].
365. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the  $p\bar{p} \rightarrow t\bar{t}$  production cross-section and the top quark mass at  $\sqrt{s} = 1.96\text{-TeV}$  in the all-hadronic decay mode,” Phys. Rev. D **76** (2007) 072009 [arXiv:0706.3790 [hep-ex]].
366. T. Aaltonen *et al.* [CDF Collaboration], “Search for New Particles Leading to  $Z$ + jets Final States in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96\text{-TeV}$ ,” Phys. Rev. D **76** (2007) 072006 [arXiv:0706.3264 [hep-ex]].
367. F. Happacher, “Status Of Observed And Predicted B Anti-B Production At The Tevatron,” <http://www.slac.stanford.edu/spires/find/hep/www?irn=7190522> SPIRES entry *Prepared for 14th International Workshop on Deep Inelastic Scattering (DIS 2006), Tsukuba, Japan, 20-24 Apr 2006*
368. T. Aaltonen *et al.* [CDF Collaboration], “Limits on Anomalous Triple Gauge Couplings in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96\text{-TeV}$ ,” Phys. Rev. D **76** (2007) 111103 [arXiv:0705.2247 [hep-ex]].
369. T. Aaltonen *et al.* [CDF Collaboration], “Measurement of the top-quark mass using missing  $E_T$  + jets events with secondary vertex  $b$ -tagging at CDF II,” Phys. Rev. D **75** (2007) 111103 [arXiv:0705.1594 [hep-ex]].

370. A. Abulencia *et al.* [CDF Collaboration], “Search for heavy, long-lived particles that decay to photons at CDF II,” *Phys. Rev. Lett.* **99** (2007) 121801 [arXiv:0704.0760 [hep-ex]].
371. A. Abulencia *et al.* [CDF Collaboration], “Polarization of  $J/\psi$  and  $\psi_{2S}$  mesons produced in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$ -TeV,” *Phys. Rev. Lett.* **99** (2007) 132001 [arXiv:0704.0638 [hep-ex]].
372. A. Abulencia *et al.* [CDF Collaboration], “Precise measurement of the top quark mass in the lepton+jets topology at CDF II,” *Phys. Rev. Lett.* **99** (2007) 182002 [arXiv:hep-ex/0703045].
373. A. Abulencia *et al.* [CDF Run II Collaboration], “Measurement of  $\sigma_{\chi_{c2}}\mathcal{B}(\chi_{c2} \rightarrow J/\psi\gamma)/\sigma_{\chi_{c1}}\mathcal{B}(\chi_{c1} \rightarrow J/\psi\gamma)$  in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$ -TeV,” *Phys. Rev. Lett.* **98** (2007) 232001 [arXiv:hep-ex/0703028].
374. A. Abulencia *et al.* [CDF Collaboration], “Inclusive search for new physics with like-sign dilepton events in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$ -TeV,” *Phys. Rev. Lett.* **98** (2007) 221803 [arXiv:hep-ex/0702051].
375. A. Abulencia *et al.* [CDF Run II Collaboration], “Measurement of  $\sigma_{p\text{ anti-}p \rightarrow Z \nu} \text{ Br}(Z \rightarrow 2\tau)$  in  $p\text{ anti-}p$  collisions at  $\sqrt{s} = 1.96$  TeV,” *Phys. Rev. D* **75** (2007) 092004.
376. A. Abulencia *et al.* [CDF Collaboration], “First Measurement of the Ratio of Central-Electron to Forward-Electron W Partial Cross Sections in  $p\text{-}p$ bar Collisions at  $\sqrt{s} = 1.96$  TeV,” *Phys. Rev. Lett.* **98** (2007) 251801 [arXiv:hep-ex/0702037].
377. A. Abulencia *et al.* [CDF Run II Collaboration], “Search for new physics in lepton + photon +  $X$  events with 929 pb<sup>(-1)</sup> of  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$ -TeV,” *Phys. Rev. D* **75** (2007) 112001 [arXiv:hep-ex/0702029].
378. A. Abulencia *et al.* [CDF Collaboration], “Observation of  $WZ$  Production,” *Phys. Rev. Lett.* **98** (2007) 161801 [arXiv:hep-ex/0702027].
379. A. Abulencia *et al.* [CDF - Run II Collaboration], “Measurement of the Inclusive Jet Cross Section using the  $k_T$  algorithm in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96$  TeV with the CDF II Detector,” *Phys. Rev. D* **75** (2007) 092006 [Erratum-ibid. *D* **75** (2007) 119901] [arXiv:hep-ex/0701051].
380. A. Abulencia *et al.* [CDF Collaboration], “Search for anomalous production of multi-lepton events in  $p\text{ anti-}p$  collisions at  $\sqrt{s} = 1.96$ -TeV,”
381. A. Abulencia *et al.* [CDF Collaboration], “Observation of  $B/s_0$  anti- $B/s_0$  oscillations,” *Phys. Rev. Lett.* **97** (2006) 242003
382. A. Abulencia *et al.* [CDF Collaboration], “Measurement of the  $\Lambda_b^0$  lifetime in  $\Lambda_b^0 \rightarrow J/\psi\Lambda^0$  in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$ -TeV,”

383. A. Abulencia *et al.* [CDF - Run II Collaboration], “Measurement of the  $t$  anti- $t$  production cross section in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.96$ -TeV in the all hadronic decay mode,” *Phys. Rev. D* **74** (2006) 072005
384. A. Abulencia *et al.* [CDF Collaboration], “Measurement of the  $t$  anti- $t$  production cross section in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.96$ -TeV using lepton + jets events with jet probability  $b$ -tagging,” *Phys. Rev. D* **74** (2006) 072006
385. A. Abulencia *et al.* [CDF Collaboration], “Observation of  $B/s_0 \rightarrow \ell^+ K^+ K^-$  and measurements of branching fractions of charmless two-body decays of  $B_0$  and  $B/s_0$  mesons in anti- $p$   $p$  collisions at  $\sqrt{s} = 1.96$ -TeV,” *Phys. Rev. Lett.* **97** (2006) 211802
386. A. Abulencia *et al.* [CDF Collaboration], “Search for excited and exotic muons in the  $\mu$   $\gamma$  decay channel in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.96$ -TeV,” *Phys. Rev. Lett.* **97** (2006) 191802
387. A. Abulencia *et al.* [CDF - Run II Collaboration], “Measurement of the  $B/s_0$  anti- $B/s_0$  oscillation frequency,” *Phys. Rev. Lett.* **97** (2006) 062003
388. A. Abulencia *et al.* [CDF Collaboration] “Measurement of the  $B^+$  production cross section in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.96$ -TeV” *Phys. Rev. D* **75**, 012010 (2007)
389. A. Abulencia *et al.* [CDF Collaboration], “Measurement of the  $t$  anti- $t$  production cross section in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.96$ -TeV,” *Phys. Rev. Lett.* **97** (2006) 082004
390. A. Abulencia *et al.* [CDF Collaboration], “Search for a neutral Higgs boson decaying to a  $W$  boson pair in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.96$ -TeV,” *Phys. Rev. Lett.* **97** (2006) 081802
391. A. Abulencia *et al.* [CDF Collaboration], “Top quark mass measurement from dilepton events at CDF II with the matrix-element method,” *Phys. Rev. D* **74** (2006) 032009
392. A. Abulencia *et al.* [CDF Collaboration], “Measurement of the  $b$  jet cross section in events with a  $Z$  boson in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.96$ -TeV,” *Phys. Rev. D* **74** (2006) 032008
393. A. Abulencia *et al.* [CDF Collaboration], “Search for new physics in lepton + photon +  $X$  events with 305-pb of  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.96$ -TeV,” *Phys. Rev. Lett.* **97** (2006) 031801
394. A. Abulencia *et al.* [CDF Collaboration], “Measurement of the ratio of branching fractions  $B(D_0 \rightarrow \ell^+ K^+ \pi^-)/B(D_0 \rightarrow \ell^- K^- \pi^+)$  using the CDF II detector,” *Phys. Rev. D* **74** (2006) 031109

395. A. Abulencia *et al.* [CDF Collaboration], “Search for large extra dimensions in the production of jets and missing transverse energy in p anti-p collisions at  $\sqrt{s} = 1.96\text{-TeV}$ ,” Phys. Rev. Lett. **97** (2006) 171802
396. A. Abulencia *et al.* [CDF Collaboration], “Measurement of the t anti-t production cross section in p anti-p collisions at  $\sqrt{s} = 1.96\text{-TeV}$  using missing  $E(t) + \text{jets}$  events with secondary vertex b-tagging,” Phys. Rev. Lett. **96** (2006) 202002
397. A. Abulencia *et al.* [CDF Collaboration], “Measurement of the B/c+ meson lifetime using B/c+  $\rightarrow$  J/psi e+ nu/e,” Phys. Rev. Lett. **97** (2006) 012002
398. A. Abulencia *et al.* [CDF Collaboration], “Search for high-mass resonances decaying to e mu in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96\text{-TeV}$ ,” Phys. Rev. Lett. **96** (2006) 211802
399. A. Abulencia *et al.* [CDF Collaboration], “Search for  $Z' \rightarrow e^+e^-$  using dielectron mass and angular distribution,” Phys. Rev. Lett. **96** (2006) 211801
400. A. Abulencia *et al.* [CDF Collaboration], “Measurement of the top quark mass using template methods on dilepton events in proton antiproton collisions at  $\sqrt{s} = 1.96\text{-TeV}$ ,” Phys. Rev. D **73** (2006) 112006
401. A. Abulencia *et al.* [CDF Collaboration], “Observation of  $B_s^0 \rightarrow \psi(2S)\phi$  and measurement of ratio of branching fractions  $B(B_s^0 \rightarrow \psi(2S)\phi)/B(B_s^0 \rightarrow J/\psi\phi)$ ,” Phys. Rev. Lett. **96** (2006) 231801
402. A. Abulencia *et al.* [CDF Collaboration], “Measurement of  $\sigma(\Lambda_b^0)/\sigma(\bar{B}^0) \times BR(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-)/BR(\text{anti-}B^0 \rightarrow D^+ \pi^-)$  in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96\text{-TeV}$ ,”
403. A. Abulencia *et al.* [CDF Collaboration], “Measurement of the dipion mass spectrum in  $X(3872) \rightarrow J/\psi\pi^+\pi^-$  decays,” Phys. Rev. Lett. **96** (2006) 102002
404. A. Abulencia *et al.* [CDF Collaboration], “A search for scalar bottom quarks from gluino decays in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96\text{-TeV}$ ,” Phys. Rev. Lett. **96** (2006) 171802
405. A. Abulencia *et al.* [CDF Collaboration], “Top quark mass measurement from dilepton events at CDF II,” Phys. Rev. Lett. **96** (2006) 152002
406. A. Abulencia *et al.* [CDF Collaboration], “Measurement of mass and width of the excited charmed meson states D10 and D2\*0,” Phys. Rev. D **73** (2006) 051104
407. A. Abulencia *et al.* [CDF Collaboration], “Search for anomalous semileptonic decay of heavy flavor hadrons produced in association with a W boson at CDF II,” Phys. Rev. D **73** (2006) 051101

408. A. Abulencia *et al.* [CDF II Collaboration], “Measurement of the inclusive jet cross section using the k(t) algorithm in p anti-p collisions at  $\sqrt{s} = 1.96\text{-TeV}$ ,” Phys. Rev. Lett. **96** (2006) 122001
409. A. Abulencia *et al.* [CDF Collaboration], “Search for H to b anti-b produced in association with W bosons in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96\text{-TeV}$ ,” Phys. Rev. Lett. **96** (2006) 081803
410. M. Goncharov *et al.*, “The timing system for the CDF electromagnetic calorimeters,” Nucl. Instrum. Meth. A **565** (2006) 538
411. A. Abulencia *et al.* [CDF Collaboration], “Search for second-generation scalar leptoquarks in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96\text{-TeV}$ ,” Phys. Rev. D **73** (2006) 051102
412. A. Abulencia *et al.* [CDF Run II Collaboration], “Measurement of the inclusive jet cross section in  $p\bar{p}$  interactions at  $\sqrt{s} = 1.96\text{-TeV}$  using a cone-based jet algorithm,” Phys. Rev. D **74** (2006) 071103
413. A. Abulencia *et al.* [CDF Collaboration], “Measurement of the top quark mass with the dynamical likelihood method using lepton plus jets events with b-tags in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96\text{-TeV}$ ,” Phys. Rev. D **73** (2006) 092002
414. G. Apollinari *et al.*, “Phenomenological study of the atypical heavy flavor production observed at the Fermilab Tevatron,” Phys. Rev. D **73** (2006) 014025
415. A. Abulencia *et al.* [CDF-Run II Collaboration], “Measurement of the helicity of W bosons in top-quark decays,” Phys. Rev. D **73** (2006) 111103
416. A. Abulencia *et al.* [CDF Collaboration], “Search for charged Higgs bosons from top quark decays in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96\text{-TeV}$ ,” Phys. Rev. Lett. **96** (2006) 042003
417. A. Abulencia *et al.* [CDF Collaboration], “A search for  $t \rightarrow tau\nu q$  in  $t\bar{t}$  production,” Phys. Lett. B **639** (2006) 172
418. A. Abulencia *et al.* [CDF Collaboration], “Precision top quark mass measurement in the lepton + jets topology in p anti-p collisions at  $\sqrt{s} = 1.96\text{-TeV}$ ,” Phys. Rev. Lett. **96** (2006) 022004
419. A. Abulencia *et al.* [CDF Collaboration], “Top quark mass measurement using the template method in the lepton + jets channel at CDF II,” Phys. Rev. D **73** (2006) 032003
420. A. Bhatti *et al.*, “Determination of the jet energy scale at the Collider Detector at Fermilab,” Nucl. Instrum. Meth. A **566** (2006) 375
421. F. Happacher, P. Giromini and F. Ptohos, “Status of the observed and predicted b barb production at the Tevatron,” Phys. Rev. D **73** (2006) 014026.

422. A. Abulencia *et al.* [CDF Collaboration], “Search for  $B/s \rightarrow \mu^+ \mu^-$  and  $B/d \rightarrow \mu^+ \mu^-$  decays in p anti-p collisions with CDF II,” arXiv:hep-ex/0508036.
423. A. Abulencia *et al.* [CDF Collaboration], “Measurement of the ratios of branching fractions  $B(B/s^0 \rightarrow D/s^- \pi^+)/B(B^0 \rightarrow D^- \pi^+)$  and  $B(B^+ \rightarrow \text{anti-}D^0 \pi^+)/B(B^0 \rightarrow D^- \pi^+)$ ,” arXiv:hep-ex/0508014.
424. A. Abulencia *et al.* [CDF Collaboration], “Search for new high mass particles decaying to lepton pairs in p anti-p collisions at  $s^{*(1/2)} = 1.96\text{-TeV}$ ,” arXiv:hep-ex/0507104.
425. D. Acosta *et al.* [CDF Collaboration], “Search for  $\Lambda/b \rightarrow p \pi$  and  $\Lambda/b \rightarrow p K$  decays in p anti-p collisions at  $s^{*(1/2)} = 1.96\text{-TeV}$ ,” arXiv:hep-ex/0507067.
426. D. Acosta *et al.* [CDF Collaboration], “Search for W and Z bosons in the reaction anti-p p  $\rightarrow 2$  jets + gamma at  $s^{*(1/2)} = 1.8\text{-TeV}$ ,” arXiv:hep-ex/0507051.
427. G. Apollinari *et al.*, “Search for narrow resonances below the Upsilon mesons,” arXiv:hep-ex/0507044.
428. G. Apollinari, I. Fiori, P. Giromini, F. Happacher, S. Miscetti, A. Parri and F. Ptohos, “Study of sequential semileptonic decays of b hadrons produced at the Tevatron,” arXiv:hep-ex/0507043.
429. D. Acosta *et al.* [CDF Collaboration], “A search for supersymmetric Higgs bosons in the di-tau decay mode in p anti-p collisions at  $s^{*(1/2)} = 1.8\text{-TeV}$ ,” arXiv:hep-ex/0506042.
430. D. Acosta *et al.* [CDF Collaboration], “Measurement of the t anti-t production cross section in p anti-p collisions at  $s^{*(1/2)} = 1.96\text{-TeV}$  using lepton plus jets events with semileptonic B decays to muons,” Phys. Rev. D **72** (2005) 032002 [arXiv:hep-ex/0506001].
431. D. Acosta *et al.* [CDF Collaboration], “Measurement of  $B(t \rightarrow W b)/B(t \rightarrow W q)$  at the Collider Detector at Fermilab,” Phys. Rev. Lett. **95** (2005) 102002 [arXiv:hep-ex/0505091].
432. D. Acosta *et al.* [CDF Collaboration], “Evidence for the exclusive decay  $B/c^+ \rightarrow J/\psi \pi^+$  and measurement of the mass of the B/c meson,” arXiv:hep-ex/0505076.
433. D. Acosta *et al.* [CDF Collaboration], “Study of jet shapes in inclusive jet production in p anti-p collisions at  $s^{*(1/2)} = 1.96\text{-TeV}$ ,” arXiv:hep-ex/0505013.
434. D. Acosta *et al.* [CDF Collaboration], “Measurement of the cross section for t anti-t production in p anti-p collisions using the kinematics of lepton + jets events,” arXiv:hep-ex/0504053.

435. D. Acosta *et al.* [CDF Collaboration], “K0(S) and Lambda0 production studies in p anti-p collisions at  $s^{**}(1/2) = 1800\text{-GeV}$  and  $630\text{-GeV}$ ,” arXiv:hep-ex/0504048.
436. D. Acosta *et al.* [CDF Collaboration], “Measurement of the azimuthal angle distribution of leptons from W boson decays as a function of the W transverse momentum in p anti-p collisions at  $s^{**}(1/2) = 1.8\text{-TeV}$ ,” arXiv:hep-ex/0504020.
437. D. Acosta *et al.* [CDF Collaboration], “Search for Higgs bosons decaying into b anti-b and produced in association with a vector boson in proton antiproton collisions at  $s^{**}(1/2) = 1.8\text{-TeV}$ ,” Phys. Rev. Lett. **95** (2005) 051801 [arXiv:hep-ex/0503039].
438. D. Acosta *et al.* [CDF Collaboration], “Search for long-lived doubly-charged Higgs bosons in p anti-p collisions at  $s^{**}(1/2) = 1.96\text{-TeV}$ ,” Phys. Rev. Lett. **95** (2005) 071801 [arXiv:hep-ex/0503004].
439. D. Acosta *et al.* [CDF Collaboration], “First evidence for  $B/s_0 \rightarrow \text{Phi Phi}$  decay and measurements of branching ratio and A(CP) for  $B^+ \rightarrow \text{Phi K}^+$ ,” arXiv:hep-ex/0502044.
440. D. Acosta *et al.* [CDF Collaboration], “Measurement of the moments of the hadronic invariant mass distribution in semileptonic B decays,” Phys. Rev. D **71** (2005) 051103 [arXiv:hep-ex/0502003].
441. D. Acosta *et al.* [CDF Collaboration], “Measurement of the  $W^+ W^-$  production cross section in p anti-p collisions at  $s^{**}(1/2) = 1.96\text{-TeV}$  using dilepton events,” Phys. Rev. Lett. **94** (2005) 211801 [arXiv:hep-ex/0501050].
442. D. Acosta *et al.* [CDF Collaboration], “Measurement of the forward-backward charge asymmetry from  $W \rightarrow e \nu$  production in p anti-p collisions at  $s^{**}(1/2) = 1.96\text{-TeV}$ ,” Phys. Rev. D **71** (2005) 051104 [arXiv:hep-ex/0501023].
443. D. Acosta *et al.* [CDF Collaboration], “Search for Z Z and Z W production in p anti-p collisions at  $s^{**}(1/2) = 1.96\text{-TeV}$ ,” Phys. Rev. D **71** (2005) 091105 [arXiv:hep-ex/0501021].
444. D. Acosta *et al.* [CDF Collaboration], “Measurement of the J/psi meson and b-hadron production cross sections in p anti-p collisions at  $s^{**}(1/2) = 1960\text{-GeV}$ ,” Phys. Rev. D **71** (2005) 032001 [arXiv:hep-ex/0412071].
445. D. Acosta *et al.* [CDF Collaboration], “Analysis of decay-time dependence of angular distributions in  $B/s_0 \rightarrow J/\psi \text{Phi}$  and  $B/d_0 \rightarrow J/\psi \text{K}^*0$  decays and measurement of the lifetime difference between B/s mass eigenstates,” Phys. Rev. Lett. **94** (2005) 101803 [arXiv:hep-ex/0412057].
446. D. Acosta *et al.* [CDF Collaboration], “Measurement of the cross section for prompt diphoton production in p anti-p collisions at  $s^{**}(1/2) = 1.96\text{-TeV}$ ,” Phys. Rev. Lett. **95** (2005) 022003 [arXiv:hep-ex/0412050].

447. D. Acosta *et al.* [CDF Collaboration], “Search for anomalous kinematics in  $t$  anti- $t$  dilepton events at CDF II,” *Phys. Rev. Lett.* **95** (2005) 022001 [arXiv:hep-ex/0412042].
448. D. Acosta *et al.* [CDF Collaboration], “Measurements of bottom anti-bottom azimuthal production correlations in proton antiproton collisions at  $s^{*(1/2)} = 1.8$ -TeV,” *Phys. Rev. D* **71** (2005) 092001 [arXiv:hep-ex/0412006].
449. D. Acosta *et al.* [CDF Collaboration], “Measurement of the W boson polarization in top decay at CDF at  $s^{*(1/2)} = 1.8$ -TeV,” *Phys. Rev. D* **71** (2005) 031101 [Erratum-*ibid.* *D* **71** (2005) 059901] [arXiv:hep-ex/0411070].
450. D. Acosta *et al.* [CDF Collaboration], “Measurement of the forward-backward charge asymmetry of electron positron pairs in p anti-p collisions at  $s^{*(1/2)} = 1.96$ -TeV,” *Phys. Rev. D* **71** (2005) 052002 [arXiv:hep-ex/0411059].
451. D. Acosta *et al.* [CDF Collaboration], “Measurement of charged particle multiplicities in gluon and quark jets in p anti-p collisions at  $s^{*(1/2)} = 1.8$ -TeV,” *Phys. Rev. Lett.* **94** (2005) 171802.
452. D. Acosta *et al.* [CDF Collaboration], “Search for scalar leptoquark pairs decaying to  $\nu$  anti- $\nu$  q anti-q in p anti-p collisions at  $s^{*(1/2)} = 1.96$ -TeV,” *Phys. Rev. D* **71** (2005) 112001 [Erratum-*ibid.* *D* **71** (2005) 119901] [arXiv:hep-ex/0410076].
453. D. Acosta *et al.* [CDF Collaboration], “Search for electroweak single top quark production in p anti-p collisions at  $s^{*(1/2)} = 1.96$ -TeV,” *Phys. Rev. D* **71** (2005) 012005 [arXiv:hep-ex/0410058].
454. D. Acosta *et al.* [CDF Collaboration], “Search for anomalous production of diphoton events with missing transverse energy at CDF and limits on gauge-mediated supersymmetry-breaking models,” *Phys. Rev. D* **71** (2005) 031104 [arXiv:hep-ex/0410053].
455. D. Acosta *et al.* [CDF Collaboration], “Measurement of the  $t$  anti- $t$  production cross section in p anti-p collisions at  $s^{*(1/2)} = 1.96$ -TeV using lepton + jets events with secondary vertex b-tagging,” *Phys. Rev. D* **71** (2005) 052003 [arXiv:hep-ex/0410041].
456. D. Acosta *et al.*, “Comparison of three-jet events in p anti-p collisions at  $s^{*(1/2)} = 1.8$ -TeV to predictions from a next-to-leading order QCD calculation,” *Phys. Rev. D* **71** (2005) 032002 [arXiv:hep-ex/0410018].
457. D. Acosta *et al.* [CDF Collaboration], “Measurement of partial widths and search for direct CP violation in  $D_0$  meson decays to  $K^- K^+$  and  $\pi^- \pi^+$ ,” *Phys. Rev. Lett.* **94** (2005) 122001 [arXiv:hep-ex/0504006].

458. D. Acosta *et al.*, “Search for excited and exotic electrons in the  $e$   $\gamma$  decay channel in  $p$  anti- $p$  collisions at  $s^{**}(1/2) = 1.96\text{-TeV}$ ,” *Phys. Rev. Lett.* **94** (2005) 101802 [arXiv:hep-ex/0410013].
459. D. Acosta *et al.* [CDF II Collaboration], “Measurement of  $W$   $\gamma$  and  $Z$   $\gamma$  production in  $p$  anti- $p$  collisions at  $s^{**}(1/2) = 1.96\text{-TeV}$ ,” *Phys. Rev. Lett.* **94** (2005) 041803 [arXiv:hep-ex/0410008].
460. D. Acosta *et al.* [CDF-II Collaboration], “Measurement of the  $t$  anti- $t$  production cross section in  $p$  anti- $p$  collisions at  $s^{**}(1/2) = 1.96\text{-TeV}$  using kinematic fitting of  $b$ -tagged lepton + jet events,” *Phys. Rev. D* **71** (2005) 072005 [arXiv:hep-ex/0409029].
461. D. Acosta *et al.* [CDF II Collaboration], “First measurements of inclusive  $W$  and  $Z$  cross sections from Run II of the Tevatron collider,” *Phys. Rev. Lett.* **94** (2005) 091803 [arXiv:hep-ex/0406078].
462. D. Acosta *et al.* [CDF Collaboration], “Search for doubly-charged Higgs bosons decaying to dileptons in  $p$  anti- $p$  collisions at  $s^{**}(1/2) = 1.96\text{-TeV}$ ,” *Phys. Rev. Lett.* **93** (2004) 221802 [arXiv:hep-ex/0406073].
463. D. Acosta *et al.* [CDF Collaboration], “Inclusive search for anomalous production of high- $p(T)$  like-sign lepton pairs in  $p$  anti- $p$  collisions at  $s^{**}(1/2) = 1.8\text{-TeV}$ ,” *Phys. Rev. Lett.* **93** (2004) 061802 [arXiv:hep-ex/0405063].
464. D. Acosta *et al.* [CDF Collaboration], “Measurement of the  $t$  anti- $t$  production cross section in  $p$  anti- $p$  collisions at  $s^{**}(1/2) = 1.96\text{-TeV}$  using dilepton events,” arXiv:hep-ex/0404036.
465. D. Acosta *et al.* [CDF Collaboration], “Direct photon cross section with conversions at CDF,” arXiv:hep-ex/0404022.
466. D. Acosta *et al.* [CDF Collaboration], “The underlying event in hard interactions at the Tevatron anti- $p$   $p$  collider,” arXiv:hep-ex/0404004.
467. D. Acosta *et al.* [CDF Collaboration], “Optimized search for single top quark production at the Fermilab Tevatron,” *Phys. Rev. D* **69**, 052003 (2004).
468. D. Acosta *et al.* [CDF Collaboration], “Search for  $B/s_0 \rightarrow \mu^+ \mu^-$  and  $B/d_0 \rightarrow \mu^+ \mu^-$  decays in  $p$  anti- $p$  collisions at  $s^{**}(1/2) = 1.96\text{-TeV}$ ,” arXiv:hep-ex/0403032.
469. D. Acosta *et al.*, “Observation of the narrow state  $X(3872) \rightarrow J/\psi \pi^+ \pi^-$  in anti- $p$   $p$  collisions at  $\sqrt{s} = 1.96\text{-TeV}$ ,” arXiv:hep-ex/0312021.
470. D. Acosta *et al.*, “Heavy flavor properties of jets produced in  $p$  anti- $p$  interactions at  $\sqrt{s} = 1.8\text{-TeV}$ ,” *Phys. Rev. D* **69**, 072004 (2004) arXiv:hep-ex/0311051.

471. D. Acosta *et al.*, “Measurement of the polar-angle distribution of leptons from W boson decay as a function of the W transverse momentum in p anti-p collisions at  $\sqrt{s} = 1.8\text{-TeV}$ ,”  
arXiv:hep-ex/0311050.
472. D. Acosta *et al.*, “Inclusive double pomeron exchange at the Fermilab Tevatron anti-p p collider,”  
arXiv:hep-ex/0311023.
473. D. Acosta *et al.*, “Measurement of the average time-integrated mixing probability of b-flavored hadrons produced at the Tevatron,”  
Phys. Rev. D **69**, 012002 (2004).
474. D. Acosta *et al.*, “Search for the flavor-changing neutral current decay  $D_0 \rightarrow \mu^+ \mu^-$  in p anti-p collisions at  $\sqrt{s} = 1.96\text{-TeV}$ ,”  
Phys. Rev. D **68**, 091101 (2003).
475. D. Acosta *et al.*, “Measurement of the mass difference  $m(D/s^+) - m(D^+)$  at CDF II,”  
Phys. Rev. D **68**, 072004 (2003).
476. D. Acosta *et al.*, “Measurement of prompt charm meson production cross sections in p anti-p collisions at  $\sqrt{s} = 1.96\text{-TeV}$ ,”  
Phys. Rev. Lett. **91**, 241804 (2003).
477. D. Acosta *et al.*, “Search For Associated Production Of Upsilon And Vector Boson In P Anti-P Collisions At  $\sqrt{s} = 1.8\text{-TeV}$ ,”  
Phys. Rev. Lett. **90**, 221803 (2003).
478. D. Acosta *et al.*, “Search for pair production of scalar top quarks in R-parity violating decay modes in p anti-p collisions at  $\sqrt{s} = 1.8\text{-TeV}$ ,”  
Phys. Rev. Lett. **92**, 051803 (2004).
479. D. Acosta *et al.*, “Central pseudorapidity gaps in events with a leading antiproton at the Fermilab Tevatron anti-p p collider,”  
Phys. Rev. Lett. **91**, 011802 (2003).
480. D. Acosta *et al.*, “Search for the supersymmetric partner of the top quark in dilepton events from p anti-p collisions at  $\sqrt{s} = 1.8\text{-TeV}$ ,”  
Phys. Rev. Lett. **90**, 251801 (2003)
481. Cross Section for forward  $J/\psi$  production in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.8$  TeV. Nov 2002, D. Acosta *et al.*, Published in Phys. Rev. D 66:092001,2002
482. Serch for a  $W'$  boson decaying to a top and bottom quark in 1.8 TeV  $p$  anti- $p$  collisions.  
Sep 2002 D. Acosta *et al.*, Submitted to Phys. Rev. Lett.

483. Search for radiative B hadron decays in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.8$  TeV.  
Aug 2002, D. Acosta *et al.*, Published in Phys. Rev. D 66:112002,2002.
484. Momentum distribuon of charged particles in jets in dijet events in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.8$  TeV.  
Jun 2002, D. Acosta *et al.*, Submitted to Phys. Rev. D.
485. Measurement of the ratio of b quark production cross section in  $p$  anti- $p$  collisions at  $\sqrt{s} = 630$  GeV and in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.8$  TeV. Jun 2002, D. Acosta *et al.*, Published in Phys. Rev. D 66:032002,2002.
486. Measurement of B meson lifetime using fully reconstructed B decays in  $p$  anti- $p$  collisions at  $\sqrt{s} = 630$  GeV and in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.8$  TeV.  
May 2002, D. Acosta *et al.*, Published in Phys. Rev. D 65:092009,2002.
487. Measurement of the  $B^+$  total cross section and  $B^+$  differential cross section in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.8$  TeV.  
Nov 2001, D. Acosta *et al.*, Published in Phys. Rev. D 65:052005,2002
488. Additional studies of the anomalous  $l + \text{Met} + 2,3$  jets events observed by CDF.  
Sep 2001, G. Apollinari, M. Barone, W.C. Carithers, T. Dorigo, I. Fiori, P. Giromini, F. Happacher, M. Kruse, S. Miscetti, A. Parri, F. Ptohos, Y. Srivastava (Duke U., Fermilab, Frascati, LBL, Berkeley, Northeastern U., Padua U., , INFN Padua), Submitted to Phys.Rev.D
489. Study of the heavy flavor content of jets produced in association with  $W$  boson in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.8$  TeV.  
Sep 2001, D. Acosta *et al.*, Published in Phys. Rev. D. D65:052007,2002
490. Measurement of the strong coupling constant from inclusive jet production at the Tevatron  $p$  anti- $p$  collider.  
Aug 2001, T. Affolder *et al.*, Submitted to Phys. Rev. Lett.
491. A study of  $B^0 \rightarrow J/\psi K(*)^0 \pi^+ \pi^-$  decays with the collider detector at Fermilab.  
Aug 2001, T. Affolder *et al.*, Submitted to Phys. Rev. Lett.
492. Search for new heavy particles in the  $W Z$  final state in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.8$  TeV.  
Aug 2001, T. Affolder *et al.*, Submitted to Phys. Rev. Lett.
493. Observation of diffractive  $J/\psi$  production at the Fermilab Tevatron.  
Jul 2001, T. Affolder *et al.*, Submitted to Phys. Rev. Lett.
494. Charged particle multiplicity in jets in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.8$  TeV.  
Jul 2001, T. Affolder *et al.*, Submitted to Phys. Rev. Lett.
495. Search for quark lepton compositeness and a heavy  $W'$  boson using the electron neutrino channel in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.8$  TeV.  
Jul 2001, T. Affolder *et al.*, Submitted to Phys. Rev. D.

496. Measurement of  $d\sigma/dM$  forward backward charge asymmetry for high mass Drell-Yan  $e^+e^-$  pairs from  $p$  anti- $p$  collisions.  
Jun 2001, T. Affolder *et al.*, Submitted to Phys. Rev. Lett.
497. Searches for new physics in events with a photon and  $b$  quark jet at CDF.  
Jun 2001, T. Affolder *et al.*, Submitted to Phys. Rev. D.
498. Cross section and heavy quark composition of photon + muon events in  $p$  anti- $p$  collisions.  
Jun 2001, T. Affolder *et al.*, Submitted to Phys. Rev. D.
499. Search for Gluinos and scalar quarks in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.8$  TeV using the missing energy plus multijets signature.  
Jun 2001, T. Affolder *et al.*, Submitted to Phys. Rev. D.
500. Production of  $\chi_{c1}$  and  $\chi_{c2}$  in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.8$  TeV.  
Jun 2001, T. Affolder *et al.*, Phys. Rev. Lett. 86:3963-3968,2001
501. Double diffraction dissociation at the Fermilab Tevatron Collider.  
May 2001, T. Affolder *et al.*, Submitted to Phys. Rev. D.
502. Search for narrow diphoton resonances and for gamma gamma +  $W/Z$  signatures in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.8$  TeV.  
May 2001, T. Affolder *et al.*, Submitted to Phys. Rev. D.
503. Measurement of the inclusive jet cross section in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.8$  TeV.  
Jan 2001, T. Affolder *et al.*, Phys. Rev. D 64:032001,2001.
504. Measurement of the  $t$  anti- $t$  production cross-section in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.8$  TeV.  
Jan 2001, T. Affolder *et al.*, Submitted to Phys. Rev. D.
505. First measurement of the ratio  $B(t \rightarrow Wb)/B(t \rightarrow Wq)$  and associated limit on the CKM element  $|V(tb)|$ .  
Dec 2000, T. Affolder *et al.*, Submitted to Phys. Rev. Lett.
506. Measurement of the two jet differential cross-section in proton anti-proton collisions at  $\sqrt{s} = 1800$  GeV.  
Dec 2000, T. Affolder *et al.*, Submitted to Phys. Rev. D.
507. Measurement of the top quark  $p_T$  distribution.  
May 2000, T. Affolder *et al.*, Submitted to Phys. Rev. Lett.
508. Search for the supersymmetric partner of the top quark in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.8$  TeV.  
Nov 2000, T. Affolder *et al.*, Submitted to Phys. Rev. D.

509. Search for neutral supersymmetric Higgs boson in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.8$  TeV.  
Oct 2000, T. Affolder *et al.*, Submitted to Phys. Rev. Lett.
510. Test of enhanced leading order QCD in  $W$  boson plus jets events from 1.8 TeV anti- $p$   $p$  collisions.  
Aug 2000, T. Affolder *et al.*, Submitted to Phys. Rev. D.
511. Measurement of the  $W$  boson mass with the Collider Detector at Fermilab.  
Jul 2000, T. Affolder *et al.*, Submitted to Phys. Rev. D.
512. Measurement of the decay amplitudes of  $B^0 \rightarrow J/\psi K^*$  and  $B_s^0 \rightarrow J\psi \Phi$  decays.  
Jul 2000, T. Affolder *et al.*, Phys. Rev. Lett. 85:4668-4673,2000.
513. Measurement of the top quark mass with the Collider Detector at Fermilab.  
Jun 2000, T. Affolder *et al.*, Phys. Rev. D 63:032003,2001.
514. Measurement of  $d\sigma/dy$  for High Mass Drell Yan  $e^+e^-$  Pairs from  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.8$  TeV.  
Jun 2000, T. Affolder *et al.*, Phys. Rev. D 63:011101,2001.
515. Dijet production by double Pomeron exchange at the Fermilab Tevatron.  
May 2000, T. Affolder *et al.*, Phys. Rev. Lett. 85:4215-4220,2000.
516. Measurement of  $J/\psi$  and  $\psi(2S)$  polarization in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.8$  TeV.  
Apr 2000, T. Affolder *et al.*, Phys. Rev. Lett. 85:2886-2891,2000.
517. Search for color singlet Technicolor particles in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.8$  TeV.  
2000, T. Affolder *et al.*, Phys. Rev. Lett. 84:1110-1115,2000.
518. Direct measurement of the  $W$  boson width in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.8$  TeV.  
Apr 2000, T. Affolder *et al.*, Phys. Rev. Lett. 85:3347-3352,2000.
519. Search for second and third generation leptoquarks including production via technicolor interactions in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.8$  TeV.  
Apr 2000, T. Affolder *et al.*, Phys. Rev. Lett. 85:2056-2061,2000.
520. Limits on Gravitino production and new processes with large missing transverse energy in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.8$  TeV.  
Mar 2000, T. Affolder *et al.*, Phys. Rev. Lett. 85:1378-1383,2000.
521. Diffractive dijets with a leading anti-proton in anti- $p$   $p$  collisions at  $\sqrt{s} = 1.8$  TeV.  
Mar 2000, T. Affolder *et al.*, Phys. Rev. Lett. 84:5043-5048,2000.

522. Search for new particles decaying to top anti-top in proton anti-proton collisions at  $\sqrt{s} = 1.8$  TeV.  
Mar 2000, T. Affolder *et al.*, Phys. Rev. Lett. 85:2062-2067,2000.
523. A measurement of the differential dijet mass cross-section in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.8$  TeV.  
Dec 1999, T. Affolder *et al.*, Phys. Rev. D 61:091101,2000.
524. Search for scalar top quark production in anti- $p$   $p$  collisions at  $\sqrt{s} = 1.8$  TeV.  
Nov 1999, T. Affolder *et al.*, Phys. Rev. Lett. 84:5273-5278,2000.
525. Observation of orbitally excited  $B$  mesons in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.8$  TeV.  
Nov 1999, T. Affolder *et al.*, Submitted to Phys. Rev. D.
526. Search for the charged Higgs boson in the decay of top quark pairs in the  $e \tau$  and  $\mu \tau$  channels at  $\sqrt{s} = 1.8$  TeV.  
Nov 1999, T. Affolder *et al.*, Phys. Rev. D 62: 012004,2000.
527. Search for scalar top and scalar bottom quarks in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.8$  TeV.  
Oct 1999, T. Affolder *et al.*, Phys. Rev. Lett. 84:5704-5709,2000.
528. Production of  $\Upsilon(1S)$  mesons from  $\chi_b$  decays in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.8$  TeV.  
Oct 1999, T. Affolder *et al.*, Phys. Rev. Lett. 84:2094-2099,2000.
529. Measurement of  $b$  quark fragmentation fractions in the production of strange and light  $B$  mesons in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.8$  TeV.  
1999, T. Affolder *et al.*, Phys. Rev. D 60: 092005,1999.
530. Search for a  $W'$  boson via the decay mode  $W' \rightarrow \mu\nu_\mu$  in 1.8 TeV  $p$  anti- $p$  collisions.  
Sep 1999, T. Affolder *et al.*, Phys. Rev. Lett. 84:5716-5721,2000.
531. Measurement of the helicity of  $W$  bosons in top quark decays.  
Sep 1999, T. Affolder *et al.*, Phys. Rev. Lett. 84:216-221,2000.
532. Search for a fourth generation quark more massive than the  $Z^0$  boson in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.8$  TeV.  
Sep 1999, T. Affolder *et al.*, Phys. Rev. Lett. 84:835-840,2000.
533. Measurement of  $b$  quark fragmentation fractions in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.8$  TeV.  
Sep 1999, T. Affolder *et al.*, Phys. Rev. Lett. 84:1663-1668,2000.
534. A measurement of  $\sin 2\beta$  from  $B \rightarrow J/\psi K_s^0$  with the CDF detector.  
1999, T. Affolder *et al.*, Phys. Rev. D 61: 072005,2000.

535. The transverse momentum and total cross section of  $e^+e^-$  pairs in the  $Z$  boson region from  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.8$  TeV.  
Aug 1999, T. Affolder *et al.*, Phys. Rev. Lett. 84:845-850,2000.
536. Observation of diffractive beauty production at the Fermilab Tevatron.  
Aug 1999, T. Affolder *et al.*, Phys. Rev. Lett. 84:232-237,2000.
537. Measurement of the  $B^0$  anti- $B^0$  oscillation frequency using  $lD^{*+}$  pairs and lepton flavor tags.  
1999, T. Affolder *et al.*, Phys. Rev. D 60: 112004,1999.
538. Search for technicolor particles in lepton plus two jets and multijets channels in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.8$  TeV.  
May 1999, T. Affolder *et al.*, Submitted to Phys. Rev. D.
539. Search for the flavor changing neutral current decays  $B^+ \rightarrow \mu^+\mu^-K^+$  and  $B^0 \rightarrow \mu^+\mu^-K^{*0}$ . May 1999, T. Affolder *et al.*, Phys. Rev. Lett. 83:3378-3383,1999.
540. Measurement of  $B^0$  anti- $B^0$  flavor oscillations using jet charge and lepton flavor tagging in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.8$  TeV.  
Mar 1999, F. Abe *et al.*, Phys. Rev. D 60: 072003,1999.
541. Measurement of the  $B^0$  anti- $B^0$  oscillation frequency using dimuon data in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.8$  TeV.  
Mar 1999, F. Abe *et al.*, Phys. Rev. D 60: 051101,1999.
542. A search for  $B^0$  anti- $B^0$  oscillations using the semileptonic decay  $B^0 \rightarrow \phi l + X\nu$ .  
Dec 1998, F. Abe *et al.*, Phys. Rev. Lett. 82:3576-3580,1999.
543. Measurement of  $B$  anti- $B$  rapidity correlation in  $p$  anti- $p$  collisions at  $\sqrt{s} = 1.8$  TeV.  
Dec 1998, F. Abe *et al.*, Phys. Rev. D 60: 032001,2000.
544. Search For R Parity Violating Supersymmetry Using Like Sign Dielectrons In  $p$  Anti- $p$  Collisions at  $\sqrt{s} = 1.8$  TeV.  
Nov, 1998, F. Abe *et al.*, Phys.Rev.Lett.83:2133-2138,1999
545. Events With A Rapidity Gap Between Jets In Anti- $p$   $p$  Collisions At  $\sqrt{s} = 630$  GeV.  
Aug 1998, F. Abe *et al.*, Phys.Rev.Lett.81:5278-5283,1998.
546. Search For Third Generation Leptoquarks From Technicolor Models In  $p$  Anti- $p$  Collisions  $\sqrt{s} = 1.8$  TeV.  
Nov 1998, F. Abe *et al.*, Phys. Rev. Lett.82:3206,1999.
547. Kinematics Of  $t$  Anti- $t$  Events At Cdf.  
Oct 1998, F. Abe *et al.*, Phys. Rev. D59:092001,1999

548. Search For A Technicolor  $\Omega_t$  Particle In Events With A Photon and a  $b$  Quark Jet At Cdf.  
Oct 1998, F. Abe *et al.*, Phys.Rev.Lett.83:3124-3129,1999.
549. Measurement Of The Top Quark Mass With The Collider Detector At Fermilab.  
Oct 1998, F. Abe *et al.*, Phys. Rev. Lett.82:271-276,1999, Erratum-ibid.82:2808-2809,1999.
550. Measurement Of  $Z^0$  And Drell-Yan Production Cross-Section Using Dimuons In Anti- $p$   $p$  Collisions at  $\sqrt{s} = 1.8$  TeV.  
Sep 1998, F. Abe *et al.*, Phys. Rev. D59:052002,1999.
551. Search For New Particles Decaying To  $b$  Anti- $b$  In  $p$  Anti- $p$  Collisions at  $\sqrt{s} = 1.8$  TeV.  
Sep 1998, F. Abe *et al.*, Phys. Rev. Lett. 82:2038-2043,1999.
552. Search For The Decays  $B_s^0, B_d^0 \rightarrow e^\pm \mu^\pm$  And Pati-Salam Leptoquarks.  
Aug 1998, F. Abe *et al.*, Phys. Rev. Lett.81: 5742-5747,1998.
553. A Measurement Of The Lepton Charge Asymmetry In  $W$  Boson Decays Produced In  $p$  Anti- $p$  Collisions.  
Sep 1998, F. Abe *et al.*, Phys. Rev. Lett.81: 5754-5759,1998.
554. Search For Higgs Bosons Produced In Association With A Vector Boson In  $p$  Anti- $p$  Collisions at  $\sqrt{s} = 1.8$  TeV.  
Aug 1998, F. Abe *et al.*, Phys. Rev. Lett.81: 5748-5753,1998.
555. Measurement Of The  $B_s^0$  Meson Lifetime Using Semileptonic Decays.  
Aug 1998, F. Abe *et al.*, Phys. Rev. D59: 032004,1999
556. Search For Second Generation Leptoquarks In The Dimuon Plus Dijet Channel Of  $p$  Anti- $p$  Collisions At  $\sqrt{s} = 1.8$  TeV.  
Jul 1998, F. Abe *et al.*, Phys. Rev. Lett.81: 4806-4811,1998.
557. Searches For New Physics In Diphoton Events In  $p$  Anti- $p$  Collisions At  $\sqrt{s} = 1.8$  TeV.  
Jun 1998, F. Abe *et al.*, Phys. Rev. D59: 092002,1999.
558. Measurement Of The  $B_d^0$  - Anti- $B_d^0$  Flavor Oscillation Frequency And Study Of Same Side Flavor Tagging Of B Mesons In  $p$  Anti- $p$  Collisions.  
Jun 1998, F. Abe *et al.*, Phys. Rev. D59: 032001,1999
559. Measurement Of The Cp Violation Parameter  $\sin 2\beta$  In  $B_d^0/\bar{B}_d^0 \rightarrow J/\Psi K_s^0$  Decays.  
Jun 1998, F. Abe *et al.*, Phys. Rev. Lett. 81: 5513-5518,1998.
560. Improved Measurement Of The  $B$  And Anti- $B^0$  Meson Lifetimes Using Semileptonic Decays.  
Jun 1998, F. Abe *et al.*, Phys. Rev. D58: 092002,1998 .

561. Observation Of The  $B_c$  Meson In  $p$  Anti- $p$  Collisions At  $\sqrt{s} = 1.8$  TeV.  
May 1998, F. Abe *et al.*, Phys. Rev. Lett. 81: 2432-2437,1998.
562. Search For Longlived Parents Of  $Z^0$  Bosons In  $p$  Anti- $p$  Collisions At  $\sqrt{s} = 1.8$  TeV.  
May 1998, F. Abe *et al.*, Phys. Rev. D58: 051102,1998.
563. Measurement Of The  $\sigma(W + \geq 1Jet)/\sigma(W)$  Cross-Section Ratio From Anti- $p$   $p$  Collisions At  $\sqrt{s} = 1.8$  TeV.  
Apr 1998, F. Abe *et al.*, Phys. Rev. Lett.81: 1367-1372,1998.
564. Observation Of The  $B_c$  Meson In  $p$  Anti- $p$  Collisions At  $\sqrt{s} = 1.8$  TeV.  
May 1998, F. Abe *et al.*, Phys. Rev. D58: 112004,1998.
565. Search For The Rare Decay  $W^\pm \rightarrow D_s^\pm \gamma$  In  $p$  Anti- $p$  Collisions At  $\sqrt{s} = 1.8$  TeV.  
Apr 1998, F. Abe *et al.*, Phys. Rev. D58: 091101,1998.
566. Search For The Rare Decay  $W^\pm \rightarrow \pi^\pm + \gamma$  In Proton - Anti-Proton Collisions At  $\sqrt{s} = 1.8$  TeV.  
Mar 1998, F. Abe *et al.*, Phys. Rev. D58: 031101,1998.
567. Observation Of  $B^+ \rightarrow \psi(2s)K^+$  And  $B^0 \rightarrow \psi(2s)K^*(892)$  Decays And Measurements Of  $B$  Meson Branching Fractions Into  $J/\psi$  And  $\psi(2s)$  Final States.  
Mar 1998, F. Abe *et al.*, Phys. Rev. D58: 072001,1998.
568. Search For Chargino - Neutralino Associated Production At The Fermilab Tevatron Collider.  
Mar 1998, F. Abe *et al.*, Phys. Rev. Lett. 80: 5275-5280,1998.
569. Searches For New Physics In Diphoton Events In  $p$  Anti- $p$  Collisions At  $\sqrt{s} = 1.8$  TeV.  
Jan 1998, F. Abe *et al.*, Phys. Rev. Lett. 81: 1791-1796,1998
570. Search For The Decays  $B_d^0 \rightarrow \mu^+ \mu^-$  And  $B_s^0 \rightarrow \mu^+ \mu^-$  In  $p$  Anti- $p$  Collisions At  $\sqrt{s} = 1.8$  TeV.  
Nov 1997, F. Abe *et al.*, Phys. Rev. D57: 3811-3816,1998.
571. Observation Of Hadronic  $W$  Decays In  $t$  Anti- $t$  Events With The Collider Detector At Fermilab.  
Nov 1997, F. Abe *et al.*, Phys. Rev. Lett. 80: 5720-5725,1998.
572. Measurement Of  $B$  Hadron Lifetimes Using  $J/\psi$  Final States At Cdf.  
Oct 1997, F. Abe *et al.*, Phys. Rev. D57: 5382-5401,1998.
573. Measurement Of The Top Quark Mass And  $t$  Anti- $t$  Production Cross-Section From Dilepton Events At The Collider Detector At Fermilab.  
Sep 1997, F. Abe *et al.*, Phys. Rev. Lett. 80:2779-2784,1998.

574. Measurement Of The Top Quark Mass.  
Oct 1997, F. Abe *et al.*, Phys. Rev. Lett. 80: 2767-2772,1998.
575. Measurement Of The  $t$  Anti- $t$  Production Cross-Section In  $p$  Anti- $p$  Collisions At  $\sqrt{s} = 1.8$  TeV.  
Oct 1997, F. Abe *et al.*, Phys. Rev. Lett. 80:2773-2778,1998.
576. Measurement Of The  $B^0$  Anti- $B^0$  Oscillation Frequency In  $p$  Anti- $p$  Collisions Using  $\pi B$  Meson Charge Flavor Correlations At  $\sqrt{s} = 1.8$  TeV.  
Sep 1997, F. Abe *et al.*, Phys. Rev. Lett. 80:2057-2062,1998.
577. Search For Flavor Changing Neutral Current Decays Of The Top Quark In  $p$  Anti- $p$  Collisions At  $\sqrt{s} = 1.8$  TeV.  
Jul 1997, F. Abe *et al.*, Phys. Rev. Lett. 80: 2525-2530,1998.
578. Properties Of Jets In W Boson Events From 1.8-TeV Anti- $p$   $p$  Collisions.  
Aug 1997, F. Abe *et al.*, Phys. Rev. Lett. 79:4760-4765,1997.
579. Measurement Of The Differential Cross-Section For Events With Large Total Transverse Energy In  $p$  Anti- $p$  Collisions At  $\sqrt{s} = 1.8$  TeV.  
Aug 1997, F. Abe *et al.*, Phys. Rev. Lett. 80: 3461-3466,1998
580. Dijet Production By Color - Singlet Exchange At The Fermilab Tevatron.  
Aug 1997, F. Abe *et al.*, Phys. Rev. Lett. 80: 1156-1161,1998
581. Search For First Generation Leptoquark Pair Production In  $p$  Anti- $p$  Collisions At  $\sqrt{s} = 1.8$  TeV.  
Aug 1997, F. Abe *et al.*, Phys. Rev. Lett. 79: 4327-4332,1997.
582. Search For New Particles Decaying Into  $B$  Anti- $B$  Produced In Association With  $W$  Bosons Decaying Into  $\nu_e$  Or  $\nu_\mu$  At The Tevatron.  
Jul 1997, F. Abe *et al.*, Phys. Rev. Lett. 79: 3819-3824,1997.
583. Limits On Quark - Lepton Compositeness Scales From Dileptons Produced In 1.8-TeV  $p$  Anti- $p$  Collisions.  
May 1997, F. Abe *et al.*, Phys. Rev. Lett. 79: 2198-2203,1997.
584. Search For New Gauge Bosons Decaying Into Dileptons In Anti- $p$   $p$  Collisions At  $\sqrt{s} = 1.8$  TeV.  
May 1997, F. Abe *et al.*, Phys. Rev. Lett. 79:2192-2197,1997.
585. Properties Of Photon Plus Two Jet Events In Anti- $p$   $p$  Collisions At  $\sqrt{s} = 1.8$  TeV.  
Apr 1997, F. Abe *et al.*, Phys. Rev. D57: 67-77,1998.
586. Properties Of Six Jet Events With Large Six Jet Mass At The Fermilab Proton Anti-Proton Collider.  
Apr 1997, F. Abe *et al.*, Phys. Rev. D56: 2532-2543,1997.

587. Double Parton Scattering In Anti- $p$   $p$  Collisions At  $\sqrt{s} = 1.8$  TeV.  
Apr 1997, F. Abe *et al.*, Phys. Rev. D56: 3811-3832,1997.
588. The  $\mu\tau$  And  $e\tau$  Decays Of Top Quark Pairs Produced In  $p$  Anti- $p$  Collisions At  $\sqrt{s} = 1.8$  TeV.  
Apr 1997, F. Abe *et al.*, Phys. Rev. Lett. 79: 3585-3590,1997.
589. Measurement Of Double Parton Scattering In Anti- $p$   $p$  Collisions At  $\sqrt{s} = 1.8$  TeV.  
Apr 1997, F. Abe *et al.*, Phys. Rev. Lett. 79: 584-589,1997.
590. Search For Charged Higgs Decays Of The Top Quark Using Hadronic Decays Of The Tau Lepton.  
Apr 1997, F. Abe *et al.*, Phys. Rev. Lett. 79: 357-362,1997.
591. First Observation Of The All Hadronic Decay Of  $t$  Anti- $t$  Pairs.  
Mar 1997, F. Abe *et al.*, Phys. Rev. Lett. 79: 1992-1997,1997.
592. Measurement Of Diffractive Dijet Production At The Tevatron.  
Mar 1997, F. Abe *et al.*, Phys. Rev. Lett. 79: 2636-2641,1997.
593. Production Of  $J/\psi$  Mesons From  $\chi_c$  Meson Decays In  $p$  Anti- $p$  Collisions At  $\sqrt{s} = 1.8$  TeV.  
Feb 1997, F. Abe *et al.*, Phys. Rev. Lett. 79: 578-583,1997.
594. Search For Gluinos And Squarks At The Fermilab Tevatron Collider.  
Feb 1997, F. Abe *et al.*, Phys. Rev. D56: 1357-1362,1997.
595.  $J/\psi$  And  $\psi(2s)$  Production In  $p$  Anti- $p$  Collisions At  $\sqrt{s} = 1.8$  TeV.  
Feb 1997, F. Abe *et al.*, Phys. Rev. Lett. 79: 572-577,1997.
596. Observation Of Diffractive W Boson Production At The Tevatron.  
Jan 1997, F. Abe *et al.*, Phys. Rev. Lett. 78: 2698-2703,1997.
597. Search For New Particles Decaying To Dijets At Cdf.  
Feb 1997, F. Abe *et al.*, Phys. Rev. D55: 5263-5268,1997.
598. Search For Third Generation Leptoquarks In Anti- $p$   $p$  Collisions At  $\sqrt{s} = 1.8$  TeV.  
Dec 1996, F. Abe *et al.*, Phys. Rev. Lett. 78: 2906-2911,1997.